Product Datasheet

Brachyury Overexpression Lysate NBL1-16661

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBL1-16661

Updated 11/9/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBL1-16661



NBL1-16661

Gene ID

Notes

Brachyury Overexpression Ly	vsate
Product Information	
Unit Size	0.1 mg
Concentration	The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.
Storage	Store at -80C. Avoid freeze-thaw cycles.
Buffer	RIPA buffer
Target Molecular Weight	47.3 kDa
Product Description	
Description	Brachyury Transient Overexpression Lysate
	Expression Host: HEK293T
	Plasmid: RC218530
	Accession#: NM_003181

(10 co of	ou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control 00ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate ontains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists f 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 00mM DTT.
-----------------	---

6862

Gene Symbol	TBXT
Species	Human

Protein Tag: C-MYC/DDK

Lysate Type	Overexpression

Product Application Details	
Applications	Western Blot, SDS-Page
Recommended Dilutions	Western Blot, SDS-Page



Application Notes


This product is intended for use as a positive control in Western Blot.

Overexpression of the target protein was confirmed using an antibody to DDK (FLAG) epitope tag (NBP1-71705) present on the protein construct.

Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.

Images

Western Blot: Brachyury Overexpression Lysate (Adult Normal) [NBL1-16661] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for Brachyury.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBL1-16661

H00006862-P01-10ug Recombinant Human Brachyury GST (N-Term) Protein

233-FB-025 FGF basic/FGF2/bFGF [Unconjugated]
AF2085 FGF basic/FGF2/bFGF [Unconjugated]

314-BP-010 BMP-4 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBL1-16661

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

