Product Datasheet

TMEM14C Overexpression Lysate NBL1-17022

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBL1-17022

Updated 11/9/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBL1-17022



NBL1-17022

TMEM14C Overexpression Lysate

Product Information	
Unit Size 0.1 mg	
	of the protein of interest cannot be determined for Please contact technical support for more information.
Storage Store at -80C. Avoid free	ze-thaw cycles.
Buffer RIPA buffer	
Target Molecular Weight11.4 kDa	
Product Description	
	nt Overexpression Lysate
Expression Host: HEK29	зт
Plasmid: RC204027	
Accession#: NM_016462	2
Protein Tag: C-MYC/DD	К
(100ug), and 1 vial of 2x contains 100ug of total p	lysate (100ug), 1 vial of empty vector negative control SDS sample buffer (250ul). Each vial of cell lysate rotein (at 1 mg/ml). The 2xSDS Sample Buffer consists -HCI pH6.8, 10% Glycerol, 0.002% Bromophenol blue,
Gene ID 51522	
Gene Symbol TMEM14C	
Species Human	
polymer transfection rea volume DNA transfection collection. The cells were 150mM NaCl, 1% NP-40 PMSF and 1mM Na3VO	dishes were transiently transfected with a non-lipid gent specially designed and manufactured for large h. Transfected cells were cultured for 48hrs before e lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, , 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM 4, and then centrifuged to clarify the lysate. Protein ured by BCA protein assay kit.
Lysate Type Overexpression	
Product Application Details	
Applications Western Blot	
Recommended Dilutions Western Blot	



Application Notes	This product is intended for use as a positive control in Western Blot. Overexpression of the target protein was confirmed using an antibody to DDK (FLAG) epitope tag (<u>NBP1-71705</u>) present on the protein construct.
	Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.

imay

	Western Blot: TMEM14C Overexpression Lysate (Adult Normal) [NBL1- 17022] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for TMEM14C.	158-		
		106-	1. S.	
		79 —	and the second second	
		48 —		
		35-		
		23-	100 million - 10	
		10.10		





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBL1-17022

AF3735	Notch-2 Antibody [Unconjugated]
H00006583-A01	OCTN1/SLC22A4 Antibody
NBP1-89715	HBLD2 Antibody
NBP1-59600	SLC25A39 Antibody

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBL1-17022

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

