Product Datasheet

GW182 Antibody - BSA Free NBP1-28751

Unit Size: 100 ul

Store at 4C. Do not freeze.



Reviews: 1 Publications: 12

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-28751

Updated 2/21/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP1-28751



NBP1-28751

GW182 Antibody - BSA Free

Product Information		
Unit Size	100 ul	
Concentration	1.0 mg/ml	
Storage	Store at 4C. Do not freeze.	
Clonality	Polyclonal	
Preservative	0.09% Sodium Azide	
Isotype	IgG	
Purity	Immunogen affinity purified	
Buffer	Tris-Citrate/Phosphate (pH 7.0 - 8.0)	
Product Description		
Host	Rabbit	
Gene ID	27327	
Gene Symbol	TNRC6A	
Species	Human, Mouse	
Reactivity Notes	Mouse reactivity reported in (PMID: 29935344).	
Marker	P/GW Body Marker	
Immunogen	The immunogen recognized by this antibody maps to a region between residue 575 and 625 of human GW182/glycine-tryptophan protein of 182 kDa (NP_055309.2).	
Product Application Details		
Applications	Western Blot, Immunoprecipitation, Knockdown Validated	
Recommended Dilutions	Western Blot 1:2000-1:10000, Immunoprecipitation 6 ug/mg lysate, Knockdown Validated	

Images

Western Blot: GW182 Antibody [NBP1-28751] - Detection of Human GW182 by Western Blot. Samples: Whole cell lysate (50 ug) from HeLa, 293T, and Jurkat cells. Antibodies: Affinity purified rabbit anti-GW182 antibody NBP1-28751 used for WB at 0.1 ug/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.	kDa 460- 268- 238- 171- 117-
	71 - 55 - 41 - 31 - <u>50 50 50</u> HeLa 293T Jurkat







Page 3 of 5 v.20.1 Updated 2/21/2025 TBG-RNAi-CK2 **TBG-RNAi-F7** Western Blot: GW182 Antibody [NBP1-28751] - Cellular expression of
 PC3-LN4
 22Rv1
 PC3-LN4
 22Rv1

 T2
 T3
 T4
 T1
 T2
 T3
 T4
TBG nanocapsule uptake & RNAi-CK2 oligomer processing markers in 25-Caveoli 20 xenograft tumors(A) Expression levels for key nanocapsule entry & oligomer processing proteins were detected by immunoblot in PC3-LN4 100 Ago 1 75 & 22Rv1 cytosolic tumor lysates from the dose response studies. The 100 Ago 2 signals for four mice per group are shown, the proteins detected are 75 indicated on the right, & the size markers are indicated on the left. Two ← GW182 200 exposures are provided for GW182 in order to show detectable signals in 150 linear range in all lanes for both PC3-LN4 & 22Rv1 tumor lysates. T1, GW182 200 T2, T3, & T4 labels indicate different tumors within the treatment & (longe exposure 50 xenograft model groups. Antibody sourcing information is listed in Actin 37 Materials & Methods. Actin signal was used as the loading control. (B) Indirect immunofluorescence detection of GW182 proteins & GW bodies in PC3-LN4 tumors. Results from 3 mice treated with TNG-RNAi-CK2 & 3 mice treated with TBG-RNAi-F7 are shown. T1, T2, & T3 labels indicate different tumors. Antibody sourcing information is listed in Materials & Methods. Nuclei were counterstained with Sytox® Green. Scale bar represents 20 µm. Image collected & cropped by CiteAb from the following publication (https://www.oncotarget.com/lookup/doi/10.18632/oncotarget.11442), licensed under a CC-BY license. Not internally tested by Novus Biologicals. Western Blot: GW182 Antibody [NBP1-28751] - TNRC6A interacts with Input IP: α-Flag elF4E-my elF4E1 eIF4E2 elF4E1 elF4E2 eIF4E2. (A) Plasmids expressing proteins indicated were transfected into TNRC-Flag 6A 6B 6A 6B 6A 6A 6B HEK293T cells. At 48 h posttransfection, cells were lysed & the lysates -35 kDa WB: α-my elF4E2 elF4E1 25 kDa were immunoprecipitated (IP) in the presence of RNase A. The TNRC6B precipitates were resolved on SDS-PAGE followed by Western blotting. 170 kDa (B) A plasmid expressing Flag-tagged eIF4E1 or eIF4E2 was transfected into HEK293T cells. At 48 h posttransfection, cells were lysed & the lysates were immunoprecipitated with the anti-TNRC6A antibody or control IgG in the presence of RNase A. The precipitates were resolved on SDS-PAGE followed by Western blotting. (C) Upper: schematic representation of TNRC6A truncation mutants. Lower: A plasmid expressing the TNRC6A mutant indicated & a plasmid expressing myc-

www.novusbio.com

tagged eIF4E2 were transiently transfected into HEK293T cells. At 48 h posttransfection, cells were lysed & the lysates were immunoprecipitated with anti-Flag antibody in the presence of RNase A followed by Western blotting. (D) Bacterially expressed Flag-tagged eIF4E2 or eIF4E1 was incubated with Glutathione Sepharose 4B bound fusion protein of GST & the C-terminal domain of TNRC6A (6A-C). The precipitates were washed & resolved on SDS-PAGE followed by commassie brilliant blue staining (lower) & Western blotting (upper) Image collected & cropped by CiteAb

(https://academic.oup.com/proteincell/article/8/10/750/6765124), licensed under a CC-BY license. Not internally tested by Novus

from the following publication

Biologicals.



Western Blot: GW182 Antibody [NBP1-28751] - Downregulation of eIF4E2 increases the protein levels of endogenous IMP1. HeLa cells were transfected with siRNAs indicated. At 48 h posttransfection, cells were lysed. (A) A fraction of the lysate was subjected to SDS-PAGE followed by Western blotting. (B) The rest cell lysate was used to extract RNA, followed by RT-qPCR measurement of the RNA levels. Relative IMP1 protein levels were quantified with the Image J software & normalized with the β-actin levels. Translational efficiency was calculated as relative protein level divided by mRNA level. Fold repression was calculated as the value in the presence of the control siRNA divided by that in the presence of the targeting siRNA. Data presented are means \pm SD of three independent experiments. The P value is determined by twotailed Student's t test. ns, nonsignificant. *P < 0.05; **P < 0.01; ***P < 0.005. Ctrli, control siRNA; TNi, siRNAs targeting TNRC6A & TNRC6B; 4E2i, siRNA targeting eIF4E2; 4Ei, siRNA targeting eIF4E1 Image collected & cropped by CiteAb from the following publication (https://academic.oup.com/proteincell/article/8/10/750/6765124), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Seth P, Hsieh PN, Jamal S et al. Regulation of MicroRNA Machinery and Development by Interspecies S-Nitrosylation Cell 2019-02-21 [PMID: 30794773]

Shin E, Jin H, Suh DS et al. An alternative miRISC targets a cancer-associated coding sequence mutation in FOXL2 EMBO J 2020-11-20 [PMID: 33215742] (WB, Human)

Details:

Western blot analysis performed on KGN and COV434 cells transfected with wiltype FOXL2 and C134W plasmids

Shin E, Jin H, Suh DS et al. An alternative miRISC targeting a coding mutation site in FOXL2 links to granulosa cell tumor bioRxiv 2020-01-01 (WB, Human)

Wilczynska A, Gillen SL, Schmidt T, et al. eIF4A2 drives repression of translation at initiation by Ccr4-Not through purine-rich motifs in the 5\'UTR Genome Biol. 2019-12-02 [PMID: 31791371] (WB, Human)

Jeppesen, DK;Fenix, AM;Franklin, JL;Higginbotham, JN;Zhang, Q;Zimmerman, LJ;Liebler, DC;Ping, J;Liu, Q;Evans, R;Fissell, WH;Patton, JG;Rome, LH;Burnette, DT;Coffey, RJ; Reassessment of Exosome Composition Cell 2019-04-04 [PMID: 30951670] (WB, Human)

Fu Y, Chen L, Chen C et al. Crosstalk between alternative polyadenylation and miRNA in regulation of protein translational efficiency. Genome Res. 2018-09-18 [PMID: 30228199] (WB, Mouse)

Jung E, Seong Y, Jeon B et al. MicroRNAs of miR-17-92 cluster increase gene expression by targeting mRNAdestabilization pathways Biochim. Biophys. Acta 2018-06-20 [PMID: 29935344] (WB, Mouse)

Chen S, Gao G. MicroRNAs recruit eIF4E2 to repress translation of target mRNAs Protein Cell 2017-07-28 [PMID: 28755203] (WB, Human)

Ahmed K, Kren BT, Abedin MJ et al. CK2 targeted RNAi therapeutic delivered via malignant cell-directed tenfibgen nanocapsule: dose and molecular mechanisms of response in xenograft prostate tumors. Oncotarget. 2016-09-20 [PMID: 27557516] (IB, Mouse)

Meijer HA, Kong YW, Lu W et al. Translational repression and eIF4A2 activity are critical for miRNA mediated gene regulation. Science 2013-04-05 [PMID: 23559250] (WB, Human)

Kamenska A, Lu WT, Kubacka D et al. Human 4E-T represses translation of bound mRNAs and enhances microRNA-mediated silencing. Nucleic Acids Res 2013-12-13 [PMID: 24335285] (WB, Human)

Wagschal A, Rousset E, Basavarajaiah P et al. Microprocessor, Setx, Xrn2, and Rrp6 Co-operate to Induce Premature Termination of Transcription by RNAPII Cell 2012-09-14 [PMID: 22980978] (WB, Human)





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP1-28751

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP1-88231PEP	GW182 Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-28751

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

