

Product Datasheet

PRPS2 Antibody - BSA Free

NBP1-31435

Unit Size: 0.1 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 4

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-31435

Updated 2/21/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-31435



NBP1-31435

PRPS2 Antibody - BSA Free

Product Information

Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.01% Thimerosal
Isotype	IgG
Purity	Antigen Affinity-purified
Buffer	0.1M Tris (pH 7), 0.1M Glycine, 10% Glycerol
Target Molecular Weight	35 kDa

Product Description


Host	Rabbit
Gene ID	5634
Gene Symbol	PRPS2
Species	Human
Reactivity Notes	Xenopus laevis (96%).
Immunogen	Recombinant protein encompassing a sequence within the center region of human PRPS2. The exact sequence is proprietary.

Product Application Details


Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Knockdown Validated
Recommended Dilutions	Western Blot 1:500-1:3000, Immunohistochemistry 1:100-1:1000, Immunocytochemistry/ Immunofluorescence 1:100-1:1000, Immunohistochemistry-Paraffin 1:100-1:1000, Knockdown Validated

Images


Western Blot: PRPS2 Antibody [NBP1-31435] - Sample (30 ug of whole cell lysate) A: Hep G2 10% SDS PAGE PRPS2 antibody, antibody diluted at 1:1000.




Immunocytochemistry/Immunofluorescence: PRPS2 Antibody [NBP1-31435] - Analysis of HaLa, using NBP1-31435 at 1:200 dilution.




Immunohistochemistry: PRPS2 Antibody [NBP1-31435] - Representative immunohistochemistry images of PRPS2 protein expression from the whole TMA with low intensity (b1) in normal prostate tissue (K11, scale bar = 0.2 mm), with low intensity (b4) in prostate cancer tissue (A9, scale bar = 0.2 mm), with intermediate intensity (b3) in prostate cancer tissue (G5, scale bar = 0.2 mm), and with high intensity (b2) in prostate cancer tissue (D2, scale bar = 0.2 mm). The percentage of PRPS2 positive cells in D2 is 90%, in G5 is 70%, and in A9 is 15%. Therefore, the percentage score of the case in D7 is 3 and its total protein expression score is $3 \times 3 = 9$. The percentage score of the case in H7 is 2 and its total protein expression score is $2 \times 2 = 4$. The percentage score of the case in E4 is 1 and its total protein expression score is $1 \times 1 = 1$. Image collected and cropped by CiteAb from the following publication (jancer.org/v11p1027.htm), licensed under a CC-BY license.




Immunohistochemistry-Paraffin: PRPS2 Antibody [NBP1-31435] - Paraffin-embedded Hep3B xenograft, using antibody at 1:100 dilution.




Western Blot: PRPS2 Antibody [NBP1-31435] - The effects of PRPS2 knockdown confirmed by qRT-PCR (top) and Western blotting (bottom). Image collected and cropped by CiteAb from the following publication (jancer.org/v11p1027.htm), licensed under a CC-BY license.




Western Blot: PRPS2 Antibody [NBP1-31435] - Knockdown of PRPS2 decreased prostate cancer cell growth. (A) The effects of PRPS2 knockdown confirmed by qRT-PCR (top) & Western blotting (bottom). (B) In the CCK₈ assay, cell viability was decreased in si-PRPS2 compared with NC. (C) In the plate colony formation assay, colony formation was obviously decreased in si-PRPS2 cells transfected. ***P < 0.001. NC: negative controls. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31956349>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: PRPS2 Antibody [NBP1-31435] - Knockdown of PRPS2 inhibits xenograft tumor growth in vivo. (A) PRPS2 shRNA effectively blocked its protein expression in PC-3 cells. Cells were stably transfected with scramble (sh-Ctrl) or PRPS2 shRNA (sh-PRPS2). PRPS2 protein levels were normalized to α-Tubulin. (B) Gross observation of xenograft tumor size in NOD/SCID mice. (C & D) Silencing of PRPS2 inhibited the tumor growth, including tumor volume (P < 0.001) & weight (P = 0.028, n = 6). (E) H&E stained paraffin embedded sections obtained from xenografts of PC-3 cells. (F) Top: Immunohistochemical analysis of Ki-67 in the xenografts; Bottom: The apoptosis in tumor tissues was evaluated by TUNEL assay (×200). Graphical illustrated the quantification of Ki-67 & TUNEL positive cells percentage. ***P < 0.001. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31956349>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: PRPS2 Antibody [NBP1-31435] - PRPS2 expression in prostate cancer tissues & prostate cancer cell lines. (A) Representative immunohistochemistry images of PRPS2 protein expression from the whole TMA with low intensity (b1) in normal prostate tissue (K11, scale bar = 0.2 mm), with low intensity (b4) in prostate cancer tissue (A9, scale bar = 0.2 mm), with intermediate intensity (b3) in prostate cancer tissue (G5, scale bar = 0.2 mm), & with high intensity (b2) in prostate cancer tissue (D2, scale bar = 0.2 mm). The percentage of PRPS2 positive cells in D2 is 90%, in G5 is 70%, & in A9 is 15%. Therefore, the percentage score of the case in D7 is 3 & its total protein expression score is 3 × 3 = 9. The percentage score of the case in H7 is 2 & its total protein expression score is 2 × 2 = 4. The percentage score of the case in E4 is 1 & its total protein expression score is 1 × 1 = 1. (B) Western blotting & real-time qPCR analysis of PRPS2 protein & mRNA expression in prostate cancer cell lines & normal prostate epithelial cell lines (RWPE-1). (*P < 0.05; **P < 0.01; ***P < 0.001). PRPS2: Phosphoribosyl pyrophosphate synthetase 2; TMA: tissue microarray. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31956349>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Emmanuel Benichou, Bolaji Seffou, Selin Topçu, Ophélie Renoult, Véronique Lenoir, Julien Planchais, Caroline Bonner, Catherine Postic, Carina Prip-Buus, Claire Pecqueur, Sandra Guilmeau, Marie-Clotilde Alves-Guerra, Renaud Dentin The transcription factor ChREBP Orchestrates liver carcinogenesis by coordinating the PI3K/AKT signaling and cancer metabolism Nature Communications 2024-02-29 [PMID: 38424041]

Chen L, Zhou Q, Zhang P et al. Direct stimulation of de novo nucleotide synthesis by O-GlcNAcylation Nature chemical biology 2023-06-12 [PMID: 37308732] (WB, Human)

Liu G, Luo Y, Hou P PRPS2 Enhances Resistance to Cisplatin via Facilitating Exosomes-mediated Macrophage M2 Polarization in Non-small Cell Lung Cancer Immunological investigations 2021-07-12 [PMID: 34251965]

Qiao H, Tan X, Lv D, et al. Phosphoribosyl pyrophosphate synthetases 2 knockdown inhibits prostate cancer progression by suppressing cell cycle and inducing cell apoptosis J. Cancer 2019-12-13 [PMID: 31956349] (IF/IHC, Human)





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP1-31435

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP2-52113-0.1mg	Recombinant Human PRPS2 His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-31435

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

