Product Datasheet

LAMP-1/CD107a Antibody NBP1-49668

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-49668

Updated 1/25/2022 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP1-49668



NBP1-49668

_AMP-1/CD107a Antibody

LAMP-1/CD107a Antibody	
Product Information	
Unit Size	0.1 ml
Concentration	1.19 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Purity	Immunogen affinity purified
Buffer	PBS
Product Description	
Host	Rabbit
Gene ID	3916
Gene Symbol	LAMP1
Species	Human, Mouse, Rat (Negative)
Reactivity Notes	Human and mouse. Antibody was negative in rat.
Marker	Late Endosome Marker
Immunogen	A genomic peptide made to an internal region of the human LAMP1 protein (within residues 250-400). [Swiss-Prot P11279]
Notes	Manufactured by Genomic Antibody Technology™. GAT <u>FAQs</u>
Product Application Details	
Applications	Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry 1:100, Immunocytochemistry/Immunofluorescence 1:50-1:100, Immunohistochemistry-Paraffin 1:100
Application Notes	This LAMP1 antibody is useful for ICC/IF and IHC. Prior to immunostaining paraffin tissues, antigen retrieval with sodium citrate buffer (pH 6.0) is



recommended.

Procedures

Immunohistochemistry-Paraffin Embedded Sections protocol specific for LAMP1 Antibody (NBP1-49668) Immunohistochemistry-Paraffin Embedded Sections

Antigen Unmasking:

Bring slides to a boil in 10 mM sodium citrate buffer (pH 6.0) then maintain at a sub-boiling temperature for 10 minutes. Cool slides on bench-top for 30 minutes.

Staining:

- 1. Wash sections in deionized water three times for 5 minutes each.
- 2. Wash sections in wash buffer for 5 minutes.
- 3. Block each section with 100-400 ul blocking solution for 1 hour at room temperature.
- 4. Remove blocking solution and add 100-400 ul diluted primary antibody. Incubate overnight at 4C.
- 5. Remove antibody solution and wash sections in wash buffer three times for 5 minutes each.
- 6. Add 100-400 ul biotinylated diluted secondary antibody. Incubate 30 minutes at room temperature.
- 7. Remove secondary antibody solution and wash sections three times with wash buffer for 5 minutes each.
- 8. Add 100-400 ul Streptavidin-HRP reagent to each section and incubate for 30 minutes at room temperature.
- 9. Wash sections three times in wash buffer for 5 minutes each.
- 10. Add 100-400 ul DAB substrate to each section and monitor staining closely.
- 11. As soon as the sections develop, immerse slides in deionized water.
- 12. Counterstain sections in hematoxylin.
- 13. Wash sections in deionized water two times for 5 minutes each.
- 14. Dehydrate sections.
- 15. Mount coverslips.

Immunocytochemistry Protocol (NBP1-49668)

Immunocytochemistry Protocol

Culture cells to appropriate density in 35 mm culture dishes or 6-well plates.

- 1. Remove culture medium and add 10% formalin to the dish. Fix at room temperature for 30 minutes.
- 2. Remove the formalin and add ice cold methanol. Incubate for 5-10 minutes.
- 3. Remove methanol and add washing solution (i.e. PBS). Be sure to not let the specimen dry out. Wash three times for 10 minutes.
- 4. To block nonspecific antibody binding incubate in 10% normal goat serum from 1 hour to overnight at room temperature.
- 5. Add primary antibody at appropriate dilution and incubate at room temperature from 2 hours to overnight at room temperature.
- 6. Remove primary antibody and replace with washing solution. Wash three times for 10 minutes.
- 7. Add secondary antibody at appropriate dilution. Incubate for 1 hour at room temperature.
- 8. Remove antibody and replace with wash solution, then wash for 10 minutes. Add Hoechst 33258 to wash solution at 1:25,0000 and incubate for 10 minutes. Wash a third time for 10 minutes.
- 9. Cells can be viewed directly after washing. The plates can also be stored in PBS containing Azide covered in Parafilm (TM). Cells can also be cover-slipped using Fluoromount, with appropriate sealing.

*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-49668

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

