Product Datasheet

HLA DQA2 Antibody NBP1-52186

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-52186

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP1-52186



NBP1-52186

HLA DQA2 Antibody

Product Information Unit Size 0.1 mg Concentration 0.5 mg/ml Storage Store at -20C. Avoid freeze-thaw cycles. Clonality Polyclonal Preservative 0.02% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Host Goat Gene ID 3118 Gene Symbol HLA-DQA2 Species Human Immunogen Peptide with sequence C-QSHGPSGQYTH corresponding to internal region according to NP_064440.1. Product Application Details Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Peptide ELISA Recommended Dilutions Western Blot 0.1-0.3ug/ml, Immunohistochemistry 0.1-0.3 ug/ml, Immunohistochem	TILA DQAZ AHIBOUY	
Concentration 0.5 mg/ml Storage Store at -20C. Avoid freeze-thaw cycles. Clonality Polyclonal Preservative 0.02% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Host Goat Gene ID 3118 Gene Symbol HLA-DQA2 Species Human Immunogen Peptide with sequence C-QSHGPSGQYTH corresponding to internal region according to NP_064440.1. Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Peptide ELISA Recommended Dilutions Western Blot 0.1-0.3ug/ml, Immunohistochemistry 0.1-0.3 ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:16000 Application Notes Western Blot: Approx 26kDa band observed in Human Bone Marrow lysates (calculated MW of 28.0kDa according to NP_064440.1). Primary incubation was 1 hour	Product Information	
Storage Store at -20C. Avoid freeze-thaw cycles. Clonality Polyclonal Preservative 0.02% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Host Goat Gene ID 3118 Gene Symbol HLA-DQA2 Species Human Immunogen Peptide with sequence C-QSHGPSGQYTH corresponding to internal region according to NP_064440.1. Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Peptide ELISA Recommended Dilutions Western Blot 0.1-0.3ug/ml, Immunohistochemistry 0.1-0.3 ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:16000 Application Notes Western blot: Approx 26kDa band observed in Human Bone Marrow lysates (calculated MW of 28.0kDa according to NP_064440.1). Primary incubation was 1 hour	Unit Size	0.1 mg
Clonality Polyclonal Preservative 0.02% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Host Goat Gene ID 3118 Gene Symbol HLA-DQA2 Species Human Immunogen Peptide with sequence C-QSHGPSGQYTH corresponding to internal region according to NP_064440.1. Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Peptide ELISA Recommended Dilutions Western Blot 0.1-0.3ug/ml, Immunohistochemistry 0.1-0.3 ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:16000 Application Notes Western Blot: Approx 26kDa band observed in Human Bone Marrow lysates (calculated MW of 28.0kDa according to NP_064440.1). Primary incubation was 1 hour	Concentration	0.5 mg/ml
Preservative 0.02% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Host Goat Gene ID 3118 Gene Symbol HLA-DQA2 Species Human Immunogen Peptide with sequence C-QSHGPSGQYTH corresponding to internal region according to NP_064440.1. Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Peptide ELISA Recommended Dilutions Western Blot 0.1-0.3ug/ml, Immunohistochemistry 0.1-0.3 ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:16000 Western blot: Approx 26kDa band observed in Human Bone Marrow lysates (calculated MW of 28.0kDa according to NP_064440.1). Primary incubation was 1 hour	Storage	Store at -20C. Avoid freeze-thaw cycles.
Isotype IgG Purity Immunogen affinity purified Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Host Goat Gene ID 3118 Gene Symbol HLA-DQA2 Species Human Immunogen Peptide with sequence C-QSHGPSGQYTH corresponding to internal region according to NP_064440.1. Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Peptide ELISA Recommended Dilutions Western Blot 0.1-0.3ug/ml, Immunohistochemistry 0.1-0.3 ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:16000 Application Notes Western blot: Approx 26kDa band observed in Human Bone Marrow lysates (calculated MW of 28.0kDa according to NP_064440.1). Primary incubation was 1 hour	Clonality	Polyclonal
Purity Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Host Gene ID 3118 Gene Symbol HLA-DQA2 Species Human Immunogen Peptide with sequence C-QSHGPSGQYTH corresponding to internal region according to NP_064440.1. Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Peptide ELISA Recommended Dilutions Western Blot 0.1-0.3ug/ml, Immunohistochemistry 0.1-0.3 ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:16000 Application Notes Western blot: Approx 26kDa band observed in Human Bone Marrow lysates (calculated MW of 28.0kDa according to NP_064440.1). Primary incubation was 1 hour	Preservative	0.02% Sodium Azide
Buffer Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA Product Description Host Goat Gene ID 3118 Gene Symbol HLA-DQA2 Species Human Immunogen Peptide with sequence C-QSHGPSGQYTH corresponding to internal region according to NP_064440.1. Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Peptide ELISA Recommended Dilutions Western Blot 0.1-0.3ug/ml, Immunohistochemistry 0.1-0.3 ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:16000 Application Notes Western blot: Approx 26kDa band observed in Human Bone Marrow lysates (calculated MW of 28.0kDa according to NP_064440.1). Primary incubation was 1 hour	Isotype	IgG
Product Description Host Goat Gene ID 3118 Gene Symbol HLA-DQA2 Species Human Immunogen Peptide with sequence C-QSHGPSGQYTH corresponding to internal region according to NP_064440.1. Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Peptide ELISA Recommended Dilutions Western Blot 0.1-0.3ug/ml, Immunohistochemistry 0.1-0.3 ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:16000 Application Notes Western blot: Approx 26kDa band observed in Human Bone Marrow lysates (calculated MW of 28.0kDa according to NP_064440.1). Primary incubation was 1 hour	Purity	Immunogen affinity purified
Host Gene ID 3118 Gene Symbol HLA-DQA2 Species Human Immunogen Peptide with sequence C-QSHGPSGQYTH corresponding to internal region according to NP_064440.1. Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Peptide ELISA Recommended Dilutions Western Blot 0.1-0.3ug/ml, Immunohistochemistry 0.1-0.3 ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:16000 Application Notes Western blot: Approx 26kDa band observed in Human Bone Marrow lysates (calculated MW of 28.0kDa according to NP_064440.1). Primary incubation was 1 hour	Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA
Gene Symbol HLA-DQA2 Species Human Immunogen Peptide with sequence C-QSHGPSGQYTH corresponding to internal region according to NP_064440.1. Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Peptide ELISA Recommended Dilutions Western Blot 0.1-0.3ug/ml, Immunohistochemistry 0.1-0.3 ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:16000 Application Notes Western blot: Approx 26kDa band observed in Human Bone Marrow lysates (calculated MW of 28.0kDa according to NP_064440.1). Primary incubation was 1 hour	Product Description	
Gene Symbol HLA-DQA2 Species Human Peptide with sequence C-QSHGPSGQYTH corresponding to internal region according to NP_064440.1. Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Peptide ELISA Recommended Dilutions Western Blot 0.1-0.3ug/ml, Immunohistochemistry 0.1-0.3 ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:16000 Application Notes Western blot: Approx 26kDa band observed in Human Bone Marrow lysates (calculated MW of 28.0kDa according to NP_064440.1). Primary incubation was 1 hour	Host	Goat
Species	Gene ID	3118
Immunogen	Gene Symbol	HLA-DQA2
according to NP_064440.1. Product Application Details Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Peptide ELISA Recommended Dilutions Western Blot 0.1-0.3ug/ml, Immunohistochemistry 0.1-0.3 ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:16000 Application Notes Western blot: Approx 26kDa band observed in Human Bone Marrow lysates (calculated MW of 28.0kDa according to NP_064440.1). Primary incubation was 1 hour	Species	Human
Applications Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Peptide ELISA Recommended Dilutions Western Blot 0.1-0.3ug/ml, Immunohistochemistry 0.1-0.3 ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:16000 Application Notes Western blot: Approx 26kDa band observed in Human Bone Marrow lysates (calculated MW of 28.0kDa according to NP_064440.1). Primary incubation was 1 hour	Immunogen	
Recommended Dilutions Western Blot 0.1-0.3ug/ml, Immunohistochemistry 0.1-0.3 ug/ml, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:16000 Application Notes Western blot: Approx 26kDa band observed in Human Bone Marrow lysates (calculated MW of 28.0kDa according to NP_064440.1). Primary incubation was 1 hour	Product Application Details	
Application Notes Western blot: Approx 26kDa band observed in Human Bone Marrow lysates (calculated MW of 28.0kDa according to NP_064440.1). Primary incubation was 1 hour	Applications	
(calculated MW of 28.0kDa according to NP_064440.1). Primary incubation was 1 hour	Recommended Dilutions	
	Application Notes	(calculated MW of 28.0kDa according to NP_064440.1). Primary incubation was 1 hour



Images Western Blot: HLA DOA2 Antibody INBP

Western Blot: HLA DQA2 Antibody [NBP1-52186] - (0.3ug/ml) staining of Human Bone Marrow lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

250kDa 150kDa 100kDa 75kDa

FOL-D-


50kDa

37kDa

25kDa

20kDa

Immunohistochemistry-Paraffin: HLA DQA2 Antibody [NBP1-52186] - 3.75ug/ml staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer Ph 6, AP-staining.



Publications

Davila S, Froeling FE, Tan A et al. New genetic associations detected in a host response study to hepatitis B vaccine. Genes Immun 2010-04-01 [PMID: 20237496]





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP1-52186

HAF017 Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish

Peroxidase)]

HAF109 Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish

Peroxidase)]

NB410-28088-1mg Goat IgG Isotype Control

DY413-05 IL-13 [Biotin]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-52186

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications



