Product Datasheet

Donkey anti-Goat IgG (H+L) Secondary Antibody [Alkaline Phosphatase] (Pre-adsorbed) NBP1-74819

Unit Size: 0.5 mg

Store at 4C. After dilution, add an equal volume of glycerol to prevent loss of enzymatic activity and store at -20C. Avoid repeated freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-74819

Updated 10/30/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP1-74819



NBP1-74819

Application Notes

Donkey anti-Goat IgG (H+L) Secondary Antibody [Alkaline Phosphatase] (Pre-adsorbed)	
Product Information	
Unit Size	0.5 mg
Concentration	1.5 mg/ml
Storage	Store at 4C. After dilution, add an equal volume of glycerol to prevent loss of enzymatic activity and store at -20C. Avoid repeated freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Conjugate	Alkaline Phosphatase
Purity	Affinity purified
Buffer	30 mM Triethanolamine, pH 7.2, 5 mM Magnesium Chloride, 0.1 mM Zinc Chloride, 1 % (w/v) BSA, Protease/IgG free
Product Description	
Description	Purity > 95% based on SDS-PAGE. Donkey serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. Store undiluted liquid at 2-8C. For storage at -20C, dilute with an equal volume of glycerol to prevent loss of enzymatic activity. Prepare working dilutions prior to use and then discard.
Host	Donkey
Species	Goat
Reactivity Notes	Based on IEP, no reactivity is observed to non-immunoglobulin goat serum immunoglobulins and IgG from human, mouse, rabbit, or rat
Specificity/Sensitivity	Based on IEP, this Donkey anti-Goat IgG (H+L) Secondary Antibody [Alkaline Phosphatase] (Pre-adsorbed) heavy gamma chains on goat IgG and light chains on all goat immunoglobulins. This antibody has been pre-adsorbed against human, mouse, rabbit or rat IgG
Immunogen	This Donkey anti-Goat IgG (H+L) Secondary Antibody [Alkaline Phosphatase] (Pre-adsorbed) was developed against purified goat IgG (H&L).
Product Application Details	
Applications	Western Blot, ELISA, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 1:200-1:5000, ELISA 1:200-1:5000, Immunohistochemistry 1:500-1:5000, Immunohistochemistry-Paraffin



This conjugate is suitable for all immunoassay applications.



Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Toronto, ON M8Z 4E6

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Bio-Techne Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Frie Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Secondary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-74819

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

