

Product Datasheet

Goat anti-Canine IgG (H+L) Secondary Antibody [FITC] NBP1-75720

Unit Size: 0.5 mg

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Reviews: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-75720

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-75720



NBP1-75720

Goat anti-Canine IgG (H+L) Secondary Antibody [FITC]


Product Information	
Unit Size	0.5 mg
Concentration	1.5 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Conjugate	FITC
Purity	Affinity purified
Buffer	10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free

Product Description	
Description	Affinity purified antibody is > 95% based on SDS-PAGE. Goat serum was obtained from healthy animals of US origin and under the care of a registered veterinarian. Fluorescein-5-Isothiocyanate (FITC) Ex = 494 nm; Em = 518 nm
Host	Goat
Species	Canine
Reactivity Notes	Based on IEP, no reactivity is observed to non-immunoglobulin canine serum immunoglobulins
Specificity/Sensitivity	Based on IEP, this Goat anti-Canine IgG (H+L) Secondary Antibody [FITC] reacts with heavy gamma chains on dog IgG and light chains on all dog immunoglobulins.
Immunogen	This Goat anti-Canine IgG (H+L) Secondary Antibody [FITC] was developed against purified canine IgG (H&L).

Product Application Details	
Applications	Western Blot, ELISA, Immunoassay, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunofluorescence
Recommended Dilutions	Western Blot 1:2000 - 1:20000, ELISA 1:2000 - 1:20000, Immunohistochemistry 1:500 - 1:5000, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunoassay, Immunofluorescence
Application Notes	This conjugate is suitable for all immunoassay applications.

Images

Immunocytochemistry/Immunofluorescence: Goat anti-Canine IgG (H+L) Secondary Antibody [FITC] [NBP1-75720] - Rat liver stained with anti-nuclear antibody followed by Goat anti-Canine IgG (H+L) Secondary Antibody [FITC]. Image from verified customer review.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Secondary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-75720

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

