

Product Datasheet

TLR9 Antibody - BSA Free NBP1-76680

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 1 Publications: 4

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-76680

Updated 9/9/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-76680



NBP1-76680

TLR9 Antibody - BSA Free

Product Information

Unit Size	0.1 mg
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Peptide affinity purified
Buffer	PBS
Target Molecular Weight	97 kDa

Product Description


Description	Novus Biologicals Rabbit TLR9 Antibody - BSA Free (NBP1-76680) is a polyclonal antibody validated for use in WB, ELISA and ICC/IF. Anti-TLR9 Antibody: Cited in 4 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	54106
Gene Symbol	TLR9
Species	Human, Mouse, Rat
Specificity/Sensitivity	At least two isoforms of TLR9 are known to exist; this antibody will detect both isoforms. TLR9 antibody is predicted to not cross-react with other TLR proteins.
Immunogen	Antibody was raised against a peptide corresponding to 15 amino acids near the center of human TLR9. The immunogen is located within amino acids 840 - 890 of TLR9. Amino Acid Sequence: RGRQSGRDEDALPYD

Product Application Details


Applications	Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence
Recommended Dilutions	Western Blot 1 ug/ml, ELISA 1:100-1:2000, Immunocytochemistry/ Immunofluorescence 2-20 ug/mL

Images


Western Blot: TLR9 Antibody [NBP1-76680] - (G) Levels of TLR9 (NBP1-76680), COX4 (NB110-39115), MyD88, and NF- κ B protein by Western blot. (H) Relative expression of TLR9 and COX4 protein. Image collected and cropped by CiteAb from the following publication ([frontiersin.org/article/10.3389/fimmu.2018.01477/full](https://www.frontiersin.org/article/10.3389/fimmu.2018.01477/full)), licensed under a CC-BY license.




Immunocytochemistry/ Immunofluorescence: TLR9 Antibody - BSA Free [NBP1-76680] - Immunofluorescence of TLR9 in Mouse Spleen tissue with TLR9 antibody at 10 ug/mL.




Western Blot: TLR9 Antibody [NBP1-76680] - Analysis of TLR9 in Jurkat cell lysate with TLR9 antibody at 1 ug/mL in (A) the absence and (B) the presence of blocking peptide




Western Blot: TLR9 Antibody [NBP1-76680] - (A) Levels of COX4 (NB110-39115), TLR9 (NBP1-76680), MyD88, and NF- κ B mRNA. (B) Levels of COX4 (NB110-39115), TLR9 (NBP1-76680), MyD88, and NF- κ B protein by Western blot. Image collected and cropped by CiteAb from the following publication (frontiersin.org/article/10.3389/fimmu.2018.01477/full), licensed under a CC-BY license.




The expression levels of microtubule protein light chain 3 (LC3B), PTEN inducing putative kinase 1 (PINK1), Parkin, mitofusin 1 (Mfn1), toll-like receptor (TLR) 9, cytochrome c oxidase 4 (COX4), myeloid differentiation factor 88 (MyD88), & nuclear factor (NF)- κ B in lung tissues from animals with spontaneous breathing (CON) or mechanical ventilation at high tidal volume (HTV) with saline, antimycin A (Ama) or cyclosporine A (CsA). (A) Levels of PINK1, Parkin, & Mfn1 mRNA. (B) Levels of LC3B, PINK1, Parkin, & Mfn1 protein by Western blot. (C) Relative expression of LC3B-II/LC3B-I & PINK1 protein. (D) Relative expression of Parkin & Mfn1 protein. (E) Levels of TLR9 & COX4 mRNA. (F) Levels of MyD88 & nuclear factor- κ B (NF- κ B) mRNA. (G) Levels of TLR9, COX4, MyD88, & NF- κ B protein by Western blot. (H) Relative expression of TLR9 & COX4 protein. (I) Relative expression of MyD88 & NF- κ B protein. Fold expression for target genes was normalized to that measured for the β -actin gene. Both of these experiments were in triplicate. aP < 0.05 vs. CON group; bP < 0.05 vs. HTV group; & cP < 0.05 vs. Ama group. Image collected & cropped by CiteAb from the following publication (<https://www.frontiersin.org/article/10.3389/fimmu.2018.01477/full>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: TLR9 Antibody - BSA Free [NBP1-76680] - Levels of toll-like receptor (TLR) 9, myeloid differentiation factor 88 (MyD88), & nuclear factor (NF)- κ B in lung tissues from animals with spontaneous breathing (CON), mechanical ventilation at normal tidal volume (NTV), high tidal volume (HTV), & ODN2088 pretreatment upon HTV stimulation (HTV + ODN). aP < 0.05 vs. CON group; bP < 0.05 vs. NTV group; & cP < 0.05 vs. HTV group. Image collected & cropped by CiteAb from the following publication (<https://www.frontiersin.org/article/10.3389/fimmu.2018.01477/full>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Immunocytochemistry/ Immunofluorescence: TLR9 Antibody - BSA Free [NBP1-76680] - Immunocytochemistry of TLR9 in Jurkat cells with TLR9 antibody at 2 μ g/mL.



Publications

Jing R, Hu ZK, Lin F et al. Mitophagy-Mediated mtDNA Release Aggravates Stretching-Induced Inflammation and Lung Epithelial Cell Injury via the TLR9/MyD88/NF- κ B Pathway *Frontiers in Cell and Developmental Biology* 2020-09-04 [PMID: 33015037] (Human)

Lin JY, Jing R, Lin F et al. High Tidal Volume Induces Mitochondria Damage and Releases Mitochondrial DNA to Aggravate the Ventilator-Induced Lung Injury. *Front Immunol.* 2018-07-03 [PMID: 30018615] (WB, Rat)

Li W, Zhu S, Li J et al. Serum Amyloid A Stimulates PKR Expression and HMGB1 Release Possibly through TLR4/RAGE Receptors. *Mol Med* 2015-01-01 [PMID: 26052716]

Xu L, Wen Z, Zhou Y et al. MicroRNA-7-regulated TLR9 signaling-enhanced growth and metastatic potential of human lung cancer cells by altering the phosphoinositide-3-kinase, regulatory subunit 3/Akt pathway *Mol Biol Cell* 2013-01-01 [PMID: 23135998] (Human)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP1-76680

NBP1-76680PEP	TLR9 Antibody Blocking Peptide
NBP2-26232	CpG oligodeoxynucleotides with negative control, TLR9 ligand
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-76680

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

