Product Datasheet

GSAP Antibody - BSA Free NBP1-78400-0.1ml

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.



Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-78400

Updated 4/13/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP1-78400




NBP1-78400-0.1ml

GSAP Antibody - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS and 30% Glycerol
Product Description	
Host	Rabbit
Gene ID	54103
Gene Symbol	PION
Species	Human
Immunogen	A synthetic peptide made to an internal portion of the human GSAP protein (between residues 150-200) [UniProt A4D1B5]
Product Application Details	
Applications	Western Blot, Simple Western
Recommended Dilutions	Western Blot 1:1000, Simple Western 1:100
Application Notes	This GSAP antibody is useful for Western Blot where a band is observed ~65 kDa representing Isoform 2/3 of GSAP. In Simple Western only 10 - 15 uL of the recommended dilution is used per data point.
	See <u>Simple Western Antibody Database</u> for Simple Western validation: Tested in HepG2 lysate 0.5 mg/mL, separated by Size, antibody dilution of 1:100. Separated by Size-Wes, Sally Sue/Peggy Sue.





Publications

Jin, C;Wang, J;Wang, Y;Jia, B;Guo, X;Yang, G;Xu, P;Greengard, P;Zhou, R;Shi, Y; Modulation of amyloid precursor protein cleavage by gamma-secretase activating protein through phase separation Proceedings of the National Academy of Sciences of the United States of America [PMID: 35298330]



Procedures

Protocol specific for GSAP antibody (NBP1-78376) WB Western Blot Protocol

1. Perform SDS-PAGE on samples to be analyzed, loading 40 ug of total protein per lane.

2. Transfer proteins to membrane according to the instructions provided by the manufacturer of the membrane and transfer apparatus.

3. Stain according to standard Ponceau S procedure (or similar product) to assess transfer success, and mark molecular weight standards where appropriate.

4. Rinse the blot.

5. Block the membrane using standard blocking buffer for at least 1 hour.

6. Wash the membrane in wash buffer three times for 10 minutes each.

7. Dilute primary antibody in blocking buffer and incubate 1 hour at room temperature.

8. Wash the membrane in wash buffer three times for 10 minutes each.

9. Apply the diluted HRP conjugated secondary antibody in blocking buffer (as per manufacturers instructions) and incubate 1 hour at room temperature.

10. Wash the blot in wash buffer three times for 10 minutes each (this step can be repeated as required to reduce background).

11. Apply the detection reagent of choice in accordance with the manufacturers instructions.

Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%.

*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP1-78400-0.1ml

NBP1-42569	HepG2 Whole Cell Lysate
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-78400

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

