

Product Datasheet

Human IgM Antibody (P5E2) NBP1-78603

Unit Size: 100ug

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-78603

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-78603



NBP1-78603


Human IgM Antibody (P5E2)

Product Information	
Unit Size	100ug
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	P5E2
Preservative	0.05% Sodium Azide
Isotype	IgG1
Purity	Ion exchange chromatography
Buffer	0.01M potassium phosphate with 0.135M NaCl, 1mg/ml BSA, 2.5mM KCl
Product Description	
Host	Mouse
Species	Human
Immunogen	Purified Human Serum IgM
Product Application Details	
Applications	Western Blot, ELISA, Flow Cytometry, Immunoprecipitation, Radioimmunoassay
Recommended Dilutions	Western Blot 1:100 - 1:400, Flow Cytometry 1 ug/ml, ELISA 1 ug/ml, Immunoprecipitation 1:10 - 1:500, Radioimmunoassay 1:100 - 1:2000




Images


Western Blot: Human IgM Antibody (P5E2) [NBP1-78603] - Analysis of 1ug of purified Human IgM, Human IgA, and Human IgG and 10ul of PageRuler Prestained Protein Ladder per well.




Flow Cytometry: Human IgM Antibody (P5E2) [NBP1-78603] - Analysis of surface IgM expression was performed on Ramos (surface and cytoplasmic IgG/lambda-expressing B cells), Raji (cytoplasmic IgG/kappa-expressing B cells), and Jurkat (T cells) cell lines. Cells were stained with a mouse anti-human IgM monoclonal antibody or mouse IgG1 isotype control (black histogram), each at a concentration of 1ug/ml. After incubation of the primary antibody on ice for 1 hour, the cells were stained with a FITC-conjugated goat anti-mouse IgG Fc secondary antibody at a dilution of 1:500 for 30 minutes on ice. The representative 10,000 cells were obtained for each sample.



ELISA: Human IgM Antibody (P5E2) [NBP1-78603] - Direct ELISA analysis of various immunoglobulin (Ig) isotypes was performed by coating wells of a 96-well plate with 100ul per well of Human IgM, Human IgG, or Human IgA diluted in carbonate/bicarbonate buffer, starting at a concentration of 50ng/ml and serially diluting 2-fold to a concentration of 0.05ng/ml, overnight at 4C. Wells of the plate were washed, blocked with StartingBlock blocking buffer, and incubated with 100ul per well of a mouse anti-human IgM monoclonal antibody at a concentration of 1ug/ml for 1 hour at room temperature. The plate was washed, then incubated with 100ul per well of an HRP-conjugated goat anti-mouse IgG Fc secondary antibody at a dilution of 1:8000 for 1 hour at room temperature. Detection was performed using 1-Step Ultra TMB substrate for 5 minutes at room temperature in the dark. The reaction was stopped with 0.16M sulfuric acid, and absorbances were read on a spectrophotometer at 450-550nm.



ELISA: Human IgM Antibody (P5E2) [NBP1-78603] - Direct ELISA analysis of IgM was performed by coating wells of a 96-well plate with 100ul per well of Human IgM diluted in carbonate/bicarbonate buffer, starting at a concentration of 5ug/ml and serially diluting 2-fold to a concentration of 5ng/ml, overnight at 4C. Wells of the plate were washed, blocked with StartingBlock blocking buffer, and incubated with 100ul per well of a mouse anti-human IgM monoclonal antibody at a concentration of 1ug/ml for 1 hour at room temperature. The plate was washed, then incubated with 100ul per well of an HRP-conjugated goat anti-mouse IgG Fc secondary antibody at a dilution of 1:8000 for 1 hour at room temperature. Detection was performed using 1-Step Ultra TMB substrate for 5 minutes at room temperature in the dark. The reaction was stopped with 0.16M sulfuric acid, and absorbances were read on a spectrophotometer at 450-550nm.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP1-78603

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)
NBP1-97056-1mg	Human IgM Native Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-78603

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

