

# Product Datasheet

## Digoxin Antibody (25P1C9)

### NBP1-78617

Unit Size: 100ug

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP1-78617](http://www.novusbio.com/NBP1-78617)

Updated 12/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP1-78617](http://www.novusbio.com/reviews/destination/NBP1-78617)



**NBP1-78617**


Digoxin Antibody (25P1C9)

<b>Product Information</b>	
<b>Unit Size</b>	100ug
<b>Concentration</b>	1 mg/ml
<b>Storage</b>	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
<b>Clonality</b>	Monoclonal
<b>Clone</b>	25P1C9
<b>Preservative</b>	0.05% Sodium Azide
<b>Isotype</b>	IgG1
<b>Purity</b>	Ion exchange chromatography
<b>Buffer</b>	0.01M potassium phosphate with 0.135M NaCl, 1mg/ml BSA, 2.5mM KCl
<b>Product Description</b>	
<b>Host</b>	Mouse
<b>Species</b>	Non-species specific
<b>Reactivity Notes</b>	Chemical
<b>Immunogen</b>	Digoxin-BSA
<b>Product Application Details</b>	
<b>Applications</b>	Western Blot, Chemiluminescence Immunoassay, ELISA, Immunohistochemistry, Immunohistochemistry-Frozen, Immunoprecipitation, Radioimmunoassay
<b>Recommended Dilutions</b>	Western Blot 1 ug/ml, ELISA 1:100 - 1:2000, Immunohistochemistry, Immunoprecipitation 1:10 - 1:500, Immunohistochemistry-Frozen 1:10-1:500, Radioimmunoassay 1:100 - 1:2000, Chemiluminescence Immunoassay




## Images


**Western Blot: Digoxin Antibody (25P1C9) [NBP1-78617] - Analysis of Digoxin** was performed by loading 10ng of BSA or Digoxin-labeled BSA per well and 5ul of PageRuler Plus Prestained Protein Ladder onto a 4-20% Tris-Glycine polyacrylamide gel. Proteins were transferred to a nitrocellulose membrane using the G2 Blotter, and blocked with 5% milk in TBST for 1 hour at room temperature. BSA was detected at ~66kD using a BSA monoclonal antibody at a dilution of 1ug/ml in 5% milk in TBST overnight at 4C on a rocking platform. Digoxin-labeled BSA was detected at ~66kD using a Digoxin monoclonal antibody at a dilution of 1ug/ml in 5% milk in TBST overnight at 4C on a rocking platform. After washing, blots were probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10,000 for at least 30 minutes at room temperature. Chemiluminescent detection was performed using SuperSignal West Pico substrate.



**Chemiluminescence Immunoassay: Digoxin Antibody (25P1C9) [NBP1-78617] - Chemiluminescent detection of a Digoxin-labeled oligonucleotide** was performed by diluting a Digoxin-labeled DNA oligonucleotide (+) or an unlabeled oligonucleotide control (-) in TriTracker Loading Dye, and loading 5pmols onto a 6% DNA Retardation Gel. DNA was transferred to a Biodyne B membrane and blocked for 30 minutes at room temperature. Digoxin was detected using a Digoxin monoclonal antibody at a dilution of 1ug/ml for 30 minutes at room temperature, followed by a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10,000 for at least 30 minutes at room temperature. Chemiluminescent detection was performed using the Chemiluminescent Nucleic Acid Chemiluminescent detection Module Kit.



**ELISA: Digoxin Antibody (25P1C9) [NBP1-78617] - InDirect ELISA analysis of Digoxin** was performed by preparing a 10ug/mL solution of BSA or Digoxin-labeled BSA in 0.2M Carbonate-Bicarbonate, pH 9.4. 100uL of each solution was added to separate wells of a clear 96-well plate, and incubated overnight at 4C. Each well was blocked with SuperBlock in PBS. The plate was then washed with PBST and incubated with 100uL per well of Digoxin monoclonal antibody in triplicate at -1000 - 2pg/mL for 1 hour at room temperature. The plate was washed with PBST and incubated with 100ul per well of goat anti-mouse IgG-HRP secondary antibody in all test wells at 1:25,000 for 30 minutes at room temperature. Detection was performed using 1-Step TMB Ultra substrate for 15 minutes at room temperature. The plate was then stopped with 0.2M sulfuric acid. Absorbances were read on a spectrophotometer at 450-550nm.





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-  
techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP1-78617](http://www.novusbio.com/reviews/submit/NBP1-78617)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

