# **Product Datasheet**

# SLC38A10 Antibody - BSA Free NBP1-81193

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-81193

Updated 4/13/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP1-81193



# NBP1-81193


SLC38A10 Antibody - BSA Free

| SLC38A10 Antibody - BSA Free |  |
|------------------------------|--|
| Product Information          |  |
| Unit Size                    | 0.1 ml   |
| Concentration                | Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.   |
| Storage                      | Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.   |
| Clonality                    | Polyclonal   |
| Preservative                 | 0.02% Sodium Azide   |
| Isotype                      | IgG  |
| Purity                       | Immunogen affinity purified  |
| Buffer                       | PBS (pH 7.2) and 40% Glycerol  |
| Product Description          |  |
| Host                         | Rabbit   |
| Gene ID                      | 124565   |
| Gene Symbol                  | SLC38A10   |
| Species                      | Human  |
| Immunogen                    | This antibody was developed against Recombinant Protein corresponding to amino acids: PEDLAGEAPGGRLGEAEGLMKVEAARLSAQDPVVAVAEDGREKPKLPKEREE LEQAQIKGPVDVPGREDGKEAPEEAQLDRPGQGIAVPV  |
| Product Application Details  |  |
| Applications                 | Simple Western, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin   |
| Recommended Dilutions        | Simple Western 1:40, Immunohistochemistry 1:50 - 1:200, Immunocytochemistry/ Immunofluorescence 0.25-2 ug/ml, Immunohistochemistry-Paraffin 1:50 - 1:200   |
| Application Notes            | For IHC-Paraffin, HIER pH 6 retrieval is recommended. ICC/IF Fixation Permeabilization: Use PFA/Triton X-100.  In Simple Western only 10 - 15 uL of the recommended dilution is used per data point.  See Simple Western Antibody Database for Simple Western validation: Tested in RT-4 and U-2OS, separated by Size, antibody dilution of 1:40, apparent MW was 169 kDa. Separated by Size-Wes, Sally Sue/Peggy Sue. |




#### **Images**


Immunohistochemistry-Paraffin: SLC38A10 Antibody [NBP1-81193] - Staining in human gallbladder and skeletal muscle tissues using anti-SLC38A10 antibody. Corresponding SLC38A10 RNA-seq data are presented for the same tissues.




Immunohistochemistry-Paraffin: SLC38A10 Antibody [NBP1-81193] - Staining of human gallbladder, kidney, skeletal muscle and testis using Anti-SLC38A10 antibody NBP1-81193 (A) shows similar protein distribution across tissues to independent antibody NBP1-81194 (B).



Immunocytochemistry/Immunofluorescence: SLC38A10 Antibody [NBP1-81193] - Staining of human cell line U-2 OS shows localization to the Golgi apparatus. Antibody staining is shown in green.




Immunohistochemistry-Paraffin: SLC38A10 Antibody [NBP1-81193] - Staining of human testis.



Page 3 of 5 v.20.1 Updated 4/13/2025 Immunohistochemistry-Paraffin: SLC38A10 Antibody [NBP1-81193] -Staining of human gallbladder shows high expression. Immunohistochemistry-Paraffin: SLC38A10 Antibody [NBP1-81193] -Staining of human skeletal muscle shows low expression as expected. Immunohistochemistry-Paraffin: SLC38A10 Antibody [NBP1-81193] -Staining of human kidney. Simple Western: SLC38A10 Antibody [NBP1-81193] - Simple Western lane view shows a specific band for SLC38A10 in 0.2 mg/ml of RT-4 (left) and U-2OS (right) lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.



Simple Western: SLC38A10 Antibody [NBP1-81193] - Electropherogram image(s) of corresponding Simple Western lane view. SLC38A10 antibody was used at 1:40 dilution on RT-4 and U-2OS lysates(s).





### **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

# **Products Related to NBP1-81193**

NBP1-81193PEP SLC38A10 Recombinant Protein Antigen

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-81193

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

