## **Product Datasheet**

### Titin Antibody - BSA Free NBP1-88071

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.



#### **Publications: 4**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP1-88071

Updated 2/21/2025 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP1-88071



#### NBP1-88071

Titin Antibody - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS (pH 7.2) and 40% Glycerol
Product Description	
Host	Rabbit
Gene ID	7273
Gene Symbol	TTN
Species	Human, Rat
Reactivity Notes	Rat reactivity reported in (PMID:30337525).
Immunogen	This antibody was developed against Recombinant Protein corresponding to amino acids: GTVSTSCYLAVQVSEEFEKETTAVTEKFTTEEKRFVESRDVVMTDTSLTEEQA GPGEPAAPYFITKPVVQKLVEGGSVVFGCQVGGNPKPHVYWKKSGVPLTTGY RYKVSYNKQTGECKLVISMTFADDAGEYT
Product Application Details	
Applications	Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry 1:200 - 1:500, Immunocytochemistry/ Immunofluorescence Reported in scientific literature (PMID: 30337525), Immunohistochemistry-Paraffin 1:200 - 1:500
Application Notes	IHC-P,HIER pH6 retrieval is recommended.

#### Images

Immunohistochemistry-Paraffin: Titin Antibody [NBP1-88071] - Analysis in human skeletal muscle and liver tissues. Corresponding TTN RNA-seq data are presented for the same tissues.





TTN in Skeletal muscle

TTN in Liver

RNA [TPM] 80 160 240 320 400 RNA [TPM] 80 160 240 320 400





www.novusbio.com



Immunohistochemistry-Paraffin: Titin Antibody [NBP1-88071] - Staining of human liver shows no positivity in hepatocytes as expected.



Immunocytochemistry/ Immunofluorescence: Titin Antibody [NBP1-88071] - SMYD2 Cys13 glutathionylation or oxidation reduces myofibril integrity. a, b Monitoring the myofibril alignment in rat neonatal cardiomyocytes upon incubation of AMA (2 µg/mL) for 12 h: no expression (a) & ectopic expression of SMYD2 WT or C13S (b). Immunostainings were done by using antibodies to SMYD2, HA (green), or titin ( $\alpha$ -titin-NT, red). About 30 cells were photographed & examined for myofibril alignment or directionality by FiberFit software.34 Images represent the major myofibril structure in individual conditions. Scale bars, 10 µm. c Analysis of myofibril alignment in cardiomyocytes. Individual cell images were analyzed by the FiberFit software to determine the fiber dispersion parameter (k) values that represent the degree of fiber alignment. High k values represent the aligned networks, whereas low k values represent the disordered networks. The median values with 95% CI are shown, n = 3 independent experiments. Difference is significant by one-way ANOVA, followed by Tukey's posthoc test, \*p < 0.05, \*\*p < 0.01, \*\*\*p < 0.001, \*\*\*\*p < 0.0001 Image collected & cropped by CiteAb from the following publication (https://pubmed.ncbi.nlm.nih.gov/30337525), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Page 4 of 5 v.20.1 Updated 2/21/2025

Immunocytochemistry/ Immunofluorescence: Titin Antibody [NBP1-88071] - SMYD2 Cys13 glutathionylation induces dissociation of SMYD2 from N2A & Hsp90. a, b SMYD2 glutathionylation disrupts its interaction with Hsp90. Purified SMYD2-SH & SMYD2-SSG were incubated with GST-Hsp90 bound to glutathione beads, & eluted sample was analyzed (a). Hsp90 was co-immunoprecipitated with SMYD2 WT or C13S from HEK293 cells in response to AMA with glucose deprivation (b). c, d SMYD2 glutathionylation disrupts its interaction with N2A. Purified SMYD2-SH & SMYD2-SSG were incubated with GST-N2A bound to glutathione beads, & eluted sample was analyzed (c). FLAG-N2A was co-immunoprecipitated with SMYD2 WT or C13S in HEK293 cells in response to AMA with glucose deprivation (d). e SMYD2 subjected to glutathionylation decreases its binding with N2A. SMYD2 WT or C13S was pre-incubated with H2O2 in the absence or presence of glutathione for 15 min, then mixed with GST-N2A bound to glutathione beads for 1 h. Eluted samples were analyzed. f, g Colocalization of titin & SMYD2 decreases upon incubation of AMA in rat neonatal cardiomyocytes expressing SMYD2 WT versus C13S. Immunostainings of cardiomyocytes with antibodies to titin ( $\alpha$ -titin-NT, red), HA, or SMYD2 (green) are shown with enlarged areas for details (the red boxes) (f). Pearson's correlation coefficients were calculated to determine colocalization of titin & SMYD2 (g). About 30 cells were analyzed in individual conditions. Images represent the major colocalization pattern in individual experiments. Scale bars, 10 µm. Data represent the mean ± SD, n = 3 independent experiments Image collected & cropped by CiteAb from the following publication

(https://pubmed.ncbi.nlm.nih.gov/30337525), licensed under a CC-BY license. Not internally tested by Novus Biologicals.

www.novusbio.com



#### **Publications**

Rodriguez Garcia M, Schmeckpeper J, Landim-Vieira M et al. Disruption of Z-Disc Function Promotes Mechanical Dysfunction in Human Myocardium: Evidence for a Dual Myofilament Modulatory Role by Alpha-Actinin 2 International journal of molecular sciences 2023-09-26 [PMID: 37834023] (IHC-Fr, Human)

Munkanatta Godage DNP, VanHecke GC, Samarasinghe kTG et al. SMYD2 glutathionylation contributes to degradation of sarcomeric proteins. Nat Commun. 2018-10-18 [PMID: 30337525] (ICC/IF, Rat)

Shin SH, Huang M, Kim SH, Choi JY. Differential Protein Expression in Congenital and Acquired Cholesteatomas. PLoS One 2015-01-01 [PMID: 26335306] (IF/IHC, Human)

Seki T, Yuasa S, Kusumoto D et al. Generation and Characterization of Functional Cardiomyocytes Derived from Human T Cell-Derived Induced Pluripotent Stem Cells. PLoS One 2014-01-01 [PMID: 24465630]





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP1-88071

NBP1-88071PEP	Titin Recombinant Protein Antigen
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-88071

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

