### **Product Datasheet**

## Fucosyltransferase 8/FUT8 Overexpression Lysate NBP2-04986

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-04986

Updated 10/23/2024 v.20.1

## Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-04986



#### NBP2-04986

Fucosyltransferase 8/FUT8 Overexpression Lysate

Product Information					
Unit Size	0.1 mg				
Concentration	The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.				
Storage	Store at -80C. Avoid freeze-thaw cycles.				
Buffer	RIPA buffer				
Target Molecular Weight	66.3 kDa				
Product Description					
Description	Fucosyltransferase 8/FUT8 Transient Overexpression Lysate				
	Expression Host: HEK293T				
	Plasmid: RC223075				
	Accession#: NM_178155				
	Protein Tag: C-MYC/DDK				
	You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.				
Gene ID	2530				
Gene Symbol	FUT8				
Species	Human				
Notes	HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCI pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.				
Lysate Type	Overexpression				
Product Application Details					
Applications	Western Blot				
Recommended Dilutions	Western Blot				



# Application NotesThis product is intended for use as a positive control in Western Blot.<br/>Overexpression of the target protein was confirmed using an antibody to DDK<br/>(FLAG) epitope tag (cat# NBP1-71705) present on the protein construct.Each vial of cell lysate contains 100ug of total protein which should be sufficient<br/>for 20-50 reactions. Depending on over-expression level, antibody affinity and<br/>detection system, some lysates can go as low as 0.1 ug per load. We<br/>recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate<br/>and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before<br/>loading.

#### Images

Western Blot: FUT8 Overexpression Lysate (Adult Normal) [NBP2-04986] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for FUT8.

		 1000	
170	-		
130	-	-	
100	-	2.4	
70	-	-	
55	-	-	
40	-		
35	-		
25	-		
15	-		
10	_		





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-04986

NBP2-30689PEP	Fucosyltransferase 8/FUT8 Recombinant Protein Antigen
236-EG-200	EGF [Unconjugated]
AF5768	Fucosyltransferase 8/FUT8 Antibody [Unconjugated]
DMP900	MMP-9 [HRP]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-04986

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

