Product Datasheet

FAM177A1 Overexpression Lysate NBP2-05123

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-05123

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-05123



NBP2-05123

FAM177A1 Overexpression Lysate

Product Information	
Init Size 0.	.1 mg
	he exact concentration of the protein of interest cannot be determined for verexpression lysates. Please contact technical support for more information.
Storage St	tore at -80C. Avoid freeze-thaw cycles.
Buffer RI	IPA buffer
arget Molecular Weight25	5.9 kDa
Product Description	
E	AM177A1 Transient Overexpression Lysate
PI	lasmid: RC216009
Ad	ccession#: NM_173607
Pr	rotein Tag: C-MYC/DDK
(1 cc of	ou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control 100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate ontains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists f 4% SDS, 125mM Tris-HCI pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 00mM DTT.
Sene ID 28	83635
Sene Symbol FA	AM177A1
pecies H	uman
pc vc cc 15 Pf	EK293T cells in 10-cm dishes were transiently transfected with a non-lipid olymer transfection reagent specially designed and manufactured for large olume DNA transfection. Transfected cells were cultured for 48hrs before ollection. The cells were lysed in modified RIPA buffer (25mM Tris-HCI pH7.6, 50mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM MSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein oncentration was measured by BCA protein assay kit.
ysate Type	verexpression
Product Application Details	
Applications W	/estern Blot



Application Notes	This product is intended for use as a positive control in Western Blot. Overexpression of the target protein was confirmed using an antibody to DDK (FLAG) epitope tag (cat# NBP1-71705) present on the protein construct.
	Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.

Images





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-05123

H00283635-P01-10ug	Recombinant Human FAM177A1 GST (N-Term) Protein
1290-IL-010	IL-23A/IL-23 P19 [Unconjugated]
NBP2-33293	FAM177A1 Antibody
NB200-106	p19ARF/CDKN2A Antibody - BSA Free

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-05123

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

