Product Datasheet

Mxi1 Overexpression Lysate NBP2-05743

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-05743

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-05743



NBP2-05743

Mxi1 Overexpression Lysate


0.1 mg
The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.
Store at -80C. Avoid freeze-thaw cycles.
RIPA buffer
32.6 kDa
Mxi1 Transient Overexpression Lysate
Expression Host: HEK293T
Plasmid: RC207319
Accession#: NM_130439
Protein Tag: C-MYC/DDK
You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCI pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.
4601
MXI1
Human
HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.
Overexpression
Western Blot
Western Blot



Application NotesThis product is intended for use as a positive control in Western Blot.
Overexpression of the target protein was confirmed using an antibody to DDK
(FLAG) epitope tag (cat# NBP1-71705) present on the protein construct.Each vial of cell lysate contains 100ug of total protein which should be sufficient
for 20-50 reactions. Depending on over-expression level, antibody affinity and
detection system, some lysates can go as low as 0.1 ug per load. We
recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate
and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before
loading.

Images

Western Blot: MXI1 Overexpression Lysate (Adult Normal) [NBP2-05743] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for MXI1.







Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-05743

NB200-103	p53 Antibody (PAb 240) - BSA Free
AF4185	Mxi1 Antibody [Unconjugated]
NB600-302	c-Myc Antibody (9E10) - BSA Free
H00004601-P02-10ug	Recombinant Human Mxi1 GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-05743

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

