Product Datasheet

Asialoglycoprotein Receptor 2 Overexpression Lysate NBP2-05750

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-05750

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-05750



NBP2-05750

on of the protein of interest cannot be determined for s. Please contact technical support for more information.
reeze-thaw cycles.
ein Receptor 2 Transient Overexpression
293T
912
DDK
of lysate (100ug), 1 vial of empty vector negative control 2xSDS sample buffer (250ul). Each vial of cell lysate I protein (at 1 mg/ml). The 2xSDS Sample Buffer consists ris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue,
cm dishes were transiently transfected with a non-lipid eagent specially designed and manufactured for large ion. Transfected cells were cultured for 48hrs before ere lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM /O4, and then centrifuged to clarify the lysate. Protein assured by BCA protein assay kit.

Product Application Details	
Applications	Western Blot
Recommended Dilutions	Western Blot



Application Notes


This product is intended for use as a positive control in Western Blot.

Overexpression of the target protein was confirmed using an antibody to DDK (FLAG) epitope tag (cat# NBP1-71705) present on the protein construct.

Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.

Images

Western Blot: Asialoglycoprotein Receptor 2 Overexpression Lysate (Adult Normal) [NBP2-05750] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for Asialoglycoprotein Receptor 2.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-05750

NBP2-22697 Recombinant Human Asialoglycoprotein Receptor 2 His Protein

AF3635 GLI-2 Antibody

NBP1-85578 Asialoglycoprotein Receptor 2 Antibody

NB110-74682 Aguaporin-2 Antibody - BSA Free

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-05750

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

