### **Product Datasheet**

#### SPRTN Overexpression Lysate NBP2-06088

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-06088

Updated 10/23/2024 v.20.1

## Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-06088



#### NBP2-06088

SPRTN Overexpression Lysate

Product Information						
Unit Size	0.1 mg					
Concentration	The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.					
Storage	Store at -80C. Avoid freeze-thaw cycles.					
Buffer	RIPA buffer					
Target Molecular Weight	55 kDa					
Product Description						
Description	SPRTN Transient Overexpression Lysate					
	Expression Host: HEK293T					
	Plasmid: RC209745					
	Accession#: NM_032018					
	Protein Tag: C-MYC/DDK					
	You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCI pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.					
Gene ID	83932					
Gene Symbol	SPRTN					
Species	Human					
Notes	HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.					
Lysate Type	Overexpression					
Product Application Details						
Applications	Western Blot					
Recommended Dilutions	Western Blot					



# Application NotesThis product is intended for use as a positive control in Western Blot.<br/>Overexpression of the target protein was confirmed using an antibody to DDK<br/>(FLAG) epitope tag (cat# NBP1-71705) present on the protein construct.Each vial of cell lysate contains 100ug of total protein which should be sufficient<br/>for 20-50 reactions. Depending on over-expression level, antibody affinity and<br/>detection system, some lysates can go as low as 0.1 ug per load. We<br/>recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate<br/>and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before<br/>loading.

#### Images

Western Blot: C1orf124 Overexpression Lysate (Adult Normal) [NBP2-06088] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for C1orf124.

170	_			Ŀ		
130	—			L		
100	_			L		
70	_		-	١.		
55	_	Ľ		1		
40	_	L	-	1		
35	-	L		1		
25	-	l	2	t		
15	_					
10	_					

www.novusbio.com





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-06088

NBP1-84163PEP NB500-106	SPRTN Recombinant Protein Antigen PCNA Antibody (PC10)
NBP1-84163	SPRTN Antibody
NB100-565	FUS Antibody

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-06088

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

