Product Datasheet

PLEKHB1/PHR1 Overexpression Lysate NBP2-06551

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-06551

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-06551



NBP2-06551

PLEKHB1/PHR1 Overexpression Lysate

	•
Product Information	
Unit Size	0.1 mg
Concentration	The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.
Storage	Store at -80C. Avoid freeze-thaw cycles.
Buffer	RIPA buffer
Target Molecular Weight	27 kDa
Product Description	
Description	PLEKHB1/PHR1 Transient Overexpression Lysate
	Expression Host: HEK293T
	Plasmid: RC216627
	Accession#: NM_021200
	Protein Tag: C-MYC/DDK
	You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCI pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.
Gene ID	58473
Gene Symbol	PLEKHB1
Species	Human
Notes	HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCI pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.
Lysate Type	Overexpression
Product Application Details	
Applications	Western Blot
Recommended Dilutions	Western Blot



of cell lysate contains 100ug of total protein which should be sufficient
reactions. Depending on over-expression level, antibody affinity and system, some lysates can go as low as 0.1 ug per load. We ad starting with 5ug of cell lysate. Add an equal amount of cell lysate DS Sample buffer and boil the SDS samples for 10 minutes before

Western Blot: PLEKHB1/PHR1 Overexpression Lysate (Adult Normal) 170 — [NBP2-06551] Left-Empty vector transfected control cell lysate (HEK293) 130 — cell lysate); Right -Over-expression Lysate for PLEKHB1/PHR1. 100 — 70 — 55 — 40 — 35 — 15 — 10 — 10 — 10 —





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-06551

H00058473-P01-10ug	Recombinant Human PLEKHB1/PHR1 GST (N-Term) Protein
2914-HT-100MG	Holo-Transferrin [Unconjugated]
NBP1-28489	PLEKHB1/PHR1 Antibody
137-PS-050	P-Selectin/CD62P

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-06551

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

