## **Product Datasheet**

### FTO Overexpression Lysate NBP2-08494

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-08494

Updated 3/10/2023 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-08494



#### NBP2-08494

FTO Overexpression Lysate

Product Information		
Unit Size	0.1 mg	
Concentration	The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.	
Storage	Store at -80C. Avoid freeze-thaw cycles.	
Buffer	RIPA buffer	
Target Molecular Weight	58.1 kDa	
Product Description		
Description	FTO Transient Overexpression Lysate	
	Expression Host: HEK293T	
	Plasmid: RC215889	
	Accession#: NM_001080432	
	Protein Tag: C-MYC/DDK	
	You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.	
Gene ID	79068	
Gene Symbol	FTO	
Species	Human	
Notes	HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.This product is manufactured by and sold under license from OriGene Technologies and its use is limited solely for research purposes.	
Lysate Type	Overexpression	
Product Application Details		
Applications	Western Blot	
<b>Recommended Dilutions</b>	Western Blot	



Application Notes       This product is intended for use as a positive control in Western Blot.         Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.         Images       Western Blot: FTO Overexpression Lysate (Adult Normal) [NBP2-08494]         Left-Empty vector transfected control cell lysate (HEK293 cell lysate);       170 — 100	Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.         Images         Western Blot: FTO Overexpression Lysate (Adult Normal) [NBP2-08494]         170         170         170         170
Images Western Blot: FTO Overexpression Lysate (Adult Normal) [NBP2-08494] Left-Empty vector transfected control cell lysate (HEK293 cell lysate);	Images         Western Blot: FTO Overexpression Lysate (Adult Normal) [NBP2-08494]         Left-Empty vector transfected control cell lysate (HEK293 cell lysate);         Right -Over-expression Lysate for FTO.         100         101         102         103         104         105         105         106         107         108         109         100         100         100         100         100         100         100
Western Blot: FTO Overexpression Lysate (Adult Normal) [NBP2-08494] Left-Empty vector transfected control cell lysate (HEK293 cell lysate);	Western Blot: FTO Overexpression Lysate (Adult Normal) [NBP2-08494] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for FTO.
70 <u></u> 55 <u></u> 40 <u></u>	$   \begin{array}{c}     70 \\     55 \\     40 \\     35 \\     25 \\   \end{array} $
	25 —





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-08494

NB110-60935PEP	FTO Antibody Blocking Peptide
DLP00	Leptin/OB [HRP]
NB110-60935	FTO Antibody
DBD00	BDNF [HRP]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-08494

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

