Product Datasheet

PIGT Overexpression Lysate NBP2-10256

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-10256

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-10256



NBP2-10256

PIGT Overexpression Lysate

Concentration The exact concentration of the protein of interest cannot overexpression lysates. Please contact technical support Storage Storage Store at -80C. Avoid freeze-thaw cycles. Buffer RIPA buffer Product Description PIGT Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC230142 Accession#: NM_001184730 Protein Tag: C-MYC/DDK You will receive 1 vial of tysate (100ug), 1 vial of empty (100ug), and 1 vial of 2xSDS sample buffer (250ul). Eac contains 100ug of total protein (at 1 mg/ml). The 2xSDS of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.00 100mM DTT. Gene ID 51604 Gene Symbol PIGT Species Human Notes HEK293T cells in 10-cm dishes were transiently transfet polymer transfection. Transfected cells were culture collection. The cells were lysed in modified RIPA buffer 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inh PMSF and 1mM Na3VO4, and then centrifuged to clarify concentration was measured by BCA protein assay kit. Lysate Type Overexpression					
overexpression lysates. Please contact technical support Storage Store at -80C. Avoid freeze-thaw cycles. Buffer RIPA buffer Product Description PIGT Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC230142 Accession#: NM_001184730 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty (100ug), and 1 vial of zSDS sample buffer (250ul). Eac contains 100ug of total protein (at 1 mg/ml). The 2xSDS of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.00 100mM DTT. Gene ID 51604 Gene Symbol PIGT Species Human Notes HEK293T cells in 10-cm dishes were transiently transfeet polymer transfection reagent specially designed and ma volume DNA transfection. Transfected cells were culture collection. The cells were lysed in modified RIPA buffer 150m MACI, 1% NP-40, 1mM EDTA, 1xProteinase inh PMSF and 1mM Na3VO4, and then centrifuged to clarify concentration was measured by BCA protein assay kit. Lysate Type Overexpression	0.1 mg				
Buffer RIPA buffer Product Description PIGT Transient Overexpression Lysate Description PIGT Transient Overexpression Lysate Expression Host: HEK293T Expression Host: HEK293T Plasmid: RC230142 Accession#: NM_001184730 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vi (100ug), and 1 vial of 2xSDS sample buffer (250ul). Eac contains 100ug of total protein (at 1 mg/ml). The 2xSDS of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.00 100mM DTT. Gene ID 51604 Gene Symbol PIGT Species Human Notes HEK293T cells in 10-cm dishes were transiently transfee polymer transfection reagent specially designed and ma volume DNA transfection. Transfected cells were culture collection. The cells were lysed in modified RIPA buffer 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inh PMSF and 1mM Na3VO4, and then centrifuged to clarify concentration was measured by BCA protein assay kit. Lysate Type Overexpression	The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.				
Product Description Description PIGT Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC230142 Accession#: NM_001184730 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty 1 (100ug), and 1 vial of 2xSDS sample buffer (250ul). Eac contains 100ug of total protein (at 1 mg/ml). The 2xSDS of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.00 100mM DTT. Gene ID 51604 Gene Symbol PIGT Species Human Notes HEK293T cells in 10-cm dishes were transiently transfect polymer transfection reagent specially designed and ma volume DNA transfection. Transfected cells were culture collection. The cells were lysed in modified RIPA buffer 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inh PMSF and 1mM Na3VO4, and then centrifuged to clarify concentration was measured by BCA protein assay kit. Lysate Type Overexpression	Store at -80C. Avoid freeze-thaw cycles.				
Description PIGT Transient Overexpression Lysate Expression Host: HEK293T Expression Host: HEK293T Plasmid: RC230142 Accession#: NM_001184730 Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vi (100ug), and 1 vial of 2xSDS sample buffer (250ul). Eac contains 100ug of total protein (at 1 mg/ml). The 2xSDS of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.00 100mM DTT. Gene ID 51604 Gene Symbol PIGT Species Human Notes HEK293T cells in 10-cm dishes were transiently transfect polymer transfection. Transfected cells were culture collection. The cells were lysed in modified RIPA buffer 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inh PMSF and 1mM Na3VO4, and then centrifuged to clarify concentration was measured by BCA protein assay kit. Lysate Type Overexpression					
PIGT Transient Overexpression LysateExpression Host: HEK293TPlasmid: RC230142Accession#: NM_001184730Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty of (100ug), and 1 vial of 2xSDS sample buffer (250ul). Eac contains 100ug of total protein (at 1 mg/ml). The 2xSDS of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.00 100mM DTT.Gene ID51604Gene SymbolPIGTSpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfect polymer transfection. Transfection. Transfected cells were culture collection. The cells were lysed in modified RIPA buffer 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inh PMSF and 1mM Na3VO4, and then centrifuged to clarify concentration was measured by BCA protein assay kit.Lysate TypeOverexpression					
Plasmid: RC230142Accession#: NM_001184730Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty of (100ug), and 1 vial of 2xSDS sample buffer (250ul). Eac contains 100ug of total protein (at 1 mg/ml). The 2xSDS of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.00 100mM DTT.Gene ID51604Gene SymbolPIGTSpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfect polymer transfection reagent specially designed and ma volume DNA transfection. Transfected cells were culture collection. The cells were lysed in modified RIPA buffer 150mM Na2Cl, 1% NP-40, 1mM EDTA, 1xProteinase inh PMSF and 1mM Na3VO4, and then centrifuged to clarify concentration was measured by BCA protein assay kit.Lysate TypeOverexpression					
Accession#: NM_001184730Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty of (100ug), and 1 vial of 2xSDS sample buffer (250ul). Eac contains 100ug of total protein (at 1 mg/ml). The 2xSDS of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.00 100mM DTT.Gene ID51604Gene SymbolPIGTSpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfect polymer transfection reagent specially designed and ma volume DNA transfection. Transfected cells were culture collection. The cells were lysed in modified RIPA buffer 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inh PMSF and 1mM Na3VO4, and then centrifuged to clarify concentration was measured by BCA protein assay kit.Lysate TypeOverexpression					
Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vi (100ug), and 1 vial of 2xSDS sample buffer (250ul). Eac contains 100ug of total protein (at 1 mg/ml). The 2xSDS of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.00 100mM DTT.Gene ID51604Gene SymbolPIGTSpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfect polymer transfection reagent specially designed and ma volume DNA transfection. Transfected cells were culture collection. The cells were lysed in modified RIPA buffer 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inh PMSF and 1mM Na3VO4, and then centrifuged to clarify concentration was measured by BCA protein assay kit.Lysate TypeOverexpression					
You will receive 1 vial of lysate (100ug), 1 vial of empty v (100ug), and 1 vial of 2xSDS sample buffer (250ul). Eac contains 100ug of total protein (at 1 mg/ml). The 2xSDS of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.00 100mM DTT.Gene ID51604Gene SymbolPIGTSpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfect polymer transfection reagent specially designed and ma volume DNA transfection. Transfected cells were culture collection. The cells were lysed in modified RIPA buffer 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inh PMSF and 1mM Na3VO4, and then centrifuged to clarify concentration was measured by BCA protein assay kit.Lysate TypeOverexpression					
(100ug), and 1 vial of 2xSDS sample buffer (250ul). Eac contains 100ug of total protein (at 1 mg/ml). The 2xSDS of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.00 100mM DTT.Gene ID51604Gene SymbolPIGTSpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfect polymer transfection reagent specially designed and ma volume DNA transfection. Transfected cells were culture collection. The cells were lysed in modified RIPA buffer 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inh PMSF and 1mM Na3VO4, and then centrifuged to clarify concentration was measured by BCA protein assay kit.Lysate TypeOverexpression					
Gene SymbolPIGTSpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfect polymer transfection reagent specially designed and may volume DNA transfection. Transfected cells were cultured collection. The cells were lysed in modified RIPA buffer 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inh PMSF and 1mM Na3VO4, and then centrifuged to clarify concentration was measured by BCA protein assay kit.Lysate TypeOverexpression	h vial of cell lysate Sample Buffer consists				
SpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfect polymer transfection reagent specially designed and may volume DNA transfection. Transfected cells were culture collection. The cells were lysed in modified RIPA buffer 					
NotesHEK293T cells in 10-cm dishes were transiently transfer polymer transfection reagent specially designed and ma volume DNA transfection. Transfected cells were culture collection. The cells were lysed in modified RIPA buffer 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inh PMSF and 1mM Na3VO4, and then centrifuged to clarify concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application Details					
polymer transfection reagent specially designed and ma volume DNA transfection. Transfected cells were culture collection. The cells were lysed in modified RIPA buffer 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inh PMSF and 1mM Na3VO4, and then centrifuged to clarify concentration was measured by BCA protein assay kit.Lysate TypeOverexpressionProduct Application Details					
Product Application Details	nufactured for large d for 48hrs before 25mM Tris-HCl pH7.6, bitor cocktail mix, 1mM				
	Western Blot				
Recommended Dilutions Western Blot					
Application NotesThis product is intended for use as a positive control in V Overexpression of the target protein was confirmed usin (FLAG) epitope tag (cat# NBP1-71705) present on the p Each vial of cell lysate contains 100ug of total protein wh for 20-50 reactions. Depending on over-expression level detection system, some lysates can go as low as 0.1 ug recommend starting with 5ug of cell lysate. Add an equa and 2X SDS Sample buffer and boil the SDS samples for loading.	g an antibody to DDK rotein construct. hich should be sufficient , antibody affinity and per load. We I amount of cell lysate				



Images

Western Blot: PIGT Overexpression Lysate (Adult Normal) [NBP2-10256] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for PIGT.

170				
130	_			
100	_			
70	-			
55	-			
40	-			
35	-			
25	-			
15	-			
10	_			

www.novusbio.com





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-10256

267-N3-005	NT-3 [Unconjugated]	
DBD00	BDNF [HRP]	
NBP1-69585	PIGT Antibody	
212-GD-010	GDNF [Unconjugated]	

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-10256

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

