### **Product Datasheet**

# Smoothened Overexpression Lysate NBP2-10451

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-10451

Updated 10/23/2024 v.20.1

## Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-10451



#### NBP2-10451

Smoothened Overexpression Lysate


Unit Size    0.1 mg      Concentration    The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.      Storage    Store at -80C. Avoid freeze-thaw cycles.      Buffer    RIPA buffer      Target Molecular Weight    86.2 kDa      Product Description    Expression Host: HEK293T      Description    Plasmid: RC203451      Accession#: NM_005631    Protein Tag: C-MYC/DDK      You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl PH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.      Gene ID    6608      Gene Symbol    SMO      Species    Human      Notes    HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection. Transfected cell swere cultured for ABtrs before collection. The one inversion and annufactured for large volume DNA transfection. Transfected cells were cultured for ABtrs before collection. The one inversion were transiently transfected for ABtrs before collection. The one inversion were transfected cells were cultured for ABtrs before collection. The one inversion were transfected cells were cultured for ABtrs before collection. The one inversion wereding and manufactured for ABtrs before collection. Th	Product Information	
overexpression lysates. Please contact technical support for more information.      Storage    Store at -80C. Avoid freeze-thaw cycles.      Buffer    RIPA buffer      Target Molecular Weight    86.2 kDa      Product Description    Smoothened Transient Overexpression Lysate      Description    Expression Host: HEK293T      Plasmid: RC203451    Accession#: NM_005631      Protein Tag: C-MYC/DDK    You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/m). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.      Gene ID    6608      Gene Symbol    SMO      Species    Human      Notes    HEX293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection. Transfected cells were cultured for 48hrs before	Unit Size	0.1 mg
Buffer  RIPA buffer    Target Molecular Weight  86.2 kDa    Product Description	Concentration	
Target Molecular Weight  86.2 kDa    Product Description  Expression Host: HEK293T    Description  Smoothened Transient Overexpression Lysate    Expression Host: HEK293T  Plasmid: RC203451    Accession#: NM_005631  Protein Tag: C-MYC/DDK    You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.    Gene ID  6608    Gene Symbol  SMO    Species  Human    Notes  HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before	Storage	Store at -80C. Avoid freeze-thaw cycles.
Product Description      Description    Smoothened Transient Overexpression Lysate      Expression Host: HEK293T    Expression Host: HEK293T      Plasmid: RC203451    Accession#: NM_005631      Protein Tag: C-MYC/DDK    You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.      Gene ID    6608      Gene Symbol    SMO      Species    Human      Notes    HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before	Buffer	RIPA buffer
DescriptionSmoothened Transient Overexpression LysateExpression Host: HEK293TPlasmid: RC203451Accession#: NM_005631Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID6608Gene SymbolSMOSpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection. Transfected cells were cultured for large volume DNA transfection. Transfected cells were cultured for 48hrs before	Target Molecular Weight	86.2 kDa
Smoothened Transient Overexpression LysateExpression Host: HEK293TPlasmid: RC203451Accession#: NM_005631Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID6608Gene SymbolSMOSpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before	Product Description	
Plasmid: RC203451Accession#: NM_005631Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID6608Gene SymbolSMOSpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before	Description	
Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCI pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID6608Gene SymbolSMOSpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before		
You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID6608Gene SymbolSMOSpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before		
Gene Symbol  SMO    Species  Human    Notes  HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfected cells were cultured for 48hrs before		You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue,
Species  Human    Notes  HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfected cells were cultured for 48hrs before	Gene ID	6608
Notes    HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before	Gene Symbol	SMO
polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before	Species	Human
150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.	Notes	polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein
Lysate Type Overexpression	Lysate Type	Overexpression
Product Application Details	Product Application Details	
Applications Western Blot	Applications	Western Blot
Recommended Dilutions Western Blot	Recommended Dilutions	Western Blot



Each vial of cell lysate contains 100ug of total protein which should be sufficient
for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.

#### Images

Western Blot: Smoothened Overexpression Lysate (Adult Normal) [NBP2-10451] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for Smoothened.



www.novusbio.com





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP2-10451

H00006608-Q02-10ug	Recombinant Human Smoothened GST (N-Term) Protein
AF3690	GLI-3 Antibody
MAB10276-100	Smoothened Antibody (1015418) [Unconjugated]
464-SH-025	Sonic Hedgehog/Shh [Unconjugated]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-10451

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

