Product Datasheet

AHSA2 Antibody NBP2-12736-100ug

Unit Size: 100 ug

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-12736

Updated 1/13/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-12736




NBP2-12736-100ug

AHSA2 Antibody


100 ug 1 mg/ml Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
1 mg/ml Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw
Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw
Polyclonal
0.09% Sodium Azide
IgG
Protein A purified
PBS (pH 7.4), 50% Glycerol
Rabbit
130872
AHSA2P
Human, Mouse, Rat
Cross reacts with Aha1.
Aha2
Western Blot, Immunocytochemistry/ Immunofluorescence
Western Blot 1:100-1:2000, Immunocytochemistry/ Immunofluorescence
1 ug/ml of AHA2 Antibody was sufficient for detection of Aha2 in 20 ug of rat tissue lysate by colorimetric immunoblot analysis using Goat anti-rat IgG:HRP as the secondary Antibody.

Images

Immunocytochemistry/Immunofluorescence: AHSA2 Antibody [NBP2-12736] - =Tissue: Heat Shocked HeLa Cells. Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Rabbit Anti-AHA2 Polyclonal Antibody at 1:80 for 12 hours at 4C. Secondary Antibody: FITC Goat Anti-Rabbit (green) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Nucleus. Cytoplasm. Magnification: 100x. Heat Shocked at 42C for 1h.



Immunocytochemistry/Immunofluorescence: AHSA2 Antibody [NBP2-12736] - Tissue: Heat Shocked HeLa Cells. Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Rabbit Anti-AHA2 Polyclonal Antibody at 1:80 for 12 hours at 4C. Secondary Antibody: R-PE Goat Anti-Rabbit (yellow) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Nucleus. Cytoplasm. Magnification: 20x. Heat Shocked at 42C for 1h.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-12736

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

