Product Datasheet

Vinculin Antibody - BSA Free NBP2-20859

Unit Size: 0.1 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 3

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-20859

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-20859



NBP2-20859

Vinculin Antibody - BSA Free


Product Information	
Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.025% Proclin 300
Isotype	IgG
Purity	Antigen Affinity-purified
Buffer	PBS, 20% Glycerol
Target Molecular Weight	124 kDa
Product Description	
Description	Novus Biologicals Rabbit Vinculin Antibody - BSA Free (NBP2-20859) is a polyclonal antibody validated for use in IHC, WB and ICC/IF. Anti-Vinculin Antibody: Cited in 3 publications. All Novus Biologicals antibodies are covered by

Product Description	
Description	Novus Biologicals Rabbit Vinculin Antibody - BSA Free (NBP2-20859) is a polyclonal antibody validated for use in IHC, WB and ICC/IF. Anti-Vinculin Antibody: Cited in 3 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	7414
Gene Symbol	VCL
Species	Human, Mouse, Rat
Marker	Focal Adhesion Marker
Immunogen	Recombinant protein encompassing a sequence within the center region of human Vinculin. The exact sequence is proprietary.

Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:500-1:3000, Immunohistochemistry 1:100-1:1000, Immunocytochemistry/ Immunofluorescence 1:100-1:1000, Immunohistochemistry-Paraffin 1:100-1:1000


Images

Western Blot: Vinculin Antibody [NBP2-20859] - Mouse tissue extract (50 ug) was separated by 5% SDS-PAGE, and the membrane was blotted with Vinculin antibody [N1N3] diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (NBP2-19301) was used to detect the primary antibody.







Immunocytochemistry/Immunofluorescence: Vinculin Antibody [NBP2-20859] - Vinculin antibody detects Vinculin protein at focal adhesion by immunofluorescent analysis. Sample: HeLa cells were fixed in ice-cold MeOH for 5 min. Green: Vinculin stained by Vinculin antibody diluted at 1:500. Blue: Fluoroshield with DAPI. Scale bar= 10 um.




Immunohistochemistry-Paraffin: Vinculin Antibody [NBP2-20859] - Human hepatoma, using Vinculin antibody at 1:500 dilution. Antigen Retrieval: TrilogyTM (EDTA based) buffer, 15min




Western Blot: Vinculin Antibody [NBP2-20859] - Whole cell extracts (30 ug) was separated by 7.5% SDS-PAGE, and the membrane was blotted with Vinculin antibody diluted at 1:1000.




Western Blot: Vinculin Antibody [NBP2-20859] - Various whole cell extracts (30 ug) were separated by 7.5% SDS-PAGE, and the membrane was blotted with Vinculin antibody diluted at 1:1000. HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.



Western Blot: Vinculin Antibody [NBP2-20859] - Various whole cell extracts (30 ug) were separated by 5% SDS-PAGE, and the membrane was blotted with Vinculin antibody diluted at 1:1000. HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.



Immunohistochemistry-Paraffin: Vinculin Antibody [NBP2-20859] - Mouse testis. Vinculin antibody [N1N3] diluted at 1:500. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min.



Publications

Silver B, Gerrish K, Tokar E. Cell-free DNA as a potential biomarker of differentiation and toxicity in cardiac organoids eLife 2023-06-01 [PMID: 37261896] (Western Blot, Mouse)

Sinha BK, Murphy C, Brown SM et Al. Mechanisms of Cell Death Induced by Erastin in Human Ovarian Tumor Cells Int J Mol Sci 2024-08-29 [PMID: 39201357]

Huda N, Khambu B, Liu G et al. Senescence Connects Autophagy Deficiency to Inflammation and Tumor Progression in the Liver Cellular and molecular gastroenterology and hepatology 2022-04-08 [PMID: 35398596] (WB, Mouse)





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-20859

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

H00007414-P01-10ug Recombinant Human Vinculin GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-20859

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

