

Product Datasheet

Vitronectin Antibody NBP2-20866

Unit Size: 0.1 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-20866

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-20866



NBP2-20866

Vitronectin Antibody


Product Information	
Unit Size	0.1 ml
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.025% Proclin 300
Isotype	IgG
Purity	Antigen Affinity-purified
Buffer	PBS, 1% BSA, 20% Glycerol
Target Molecular Weight	54 kDa

Product Description	
Description	Novus Biologicals Rabbit Vitronectin Antibody (NBP2-20866) is a polyclonal antibody validated for use in IHC, WB, ELISA and ICC/IF. Anti-Vitronectin Antibody: Cited in 2 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	7448
Gene Symbol	VTN
Species	Human, Mouse
Immunogen	Recombinant protein encompassing a sequence within the center region of human Vitronectin. The exact sequence is proprietary.


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, ELISA, Immunocytochemistry/Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:500-1:3000, ELISA Assay dependent, Immunohistochemistry 1:100-1:1000, Immunocytochemistry/Immunofluorescence Assay dependent, Immunohistochemistry-Paraffin 1:100-1:1000

Images


Western Blot: Vitronectin Antibody [NBP2-20866] - Proteomics analysis of losartan-mediated reversion of early fibrotic RDEB changes. Bar graph show abundance of representative proteins normalized by losartan treatment. Mean \pm S.E.M. (LFQ intensity) of groups of individual mice (wild-type, untreated, & losartan-treated C7-hypomorphic mice). Unpaired t-test calculated significance. Abundance of Lrg1: *P-value C7-hypomorphic vs. losartan-treated C7-hypomorphic mice = 0.022; abundance of serpin f2: **P-value untreated C7-hypomorphic vs. losartan-treated C7-hypomorphic mice = 0.0084; abundance of vitronectin: **P-value untreated C7-hypomorphic vs. losartan-treated C7-hypomorphic mice = 0.0094. Western blots of skin lysates from wild-type, untreated C7-hypomorphic, & 7-week losartan-treated C7-hypomorphic mice not used for proteomics. Image collected & cropped by CiteAb from the following publication (www.onlinelibrary.wiley.com/doi/10.15252/emmm.201505061) licensed under a CC-BY license.




Immunohistochemistry-Paraffin: Vitronectin Antibody [NBP2-20866] - Vitronectin antibody detects Vitronectin protein by immunohistochemical analysis. Sample: Paraffin-embedded human lung cancer. Vitronectin stained by Vitronectin antibody diluted at 1:1000.




Immunohistochemistry-Paraffin: Vitronectin Antibody [NBP2-20866] - Human colon carcinoma, using Vitronectin antibody at 1:500 dilution. Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min.



Western Blot: Vitronectin Antibody [NBP2-20866] - HepG2 whole cell extract and conditioned medium (30 ug) were separated by 10% SDS-PAGE, and the membrane was blotted with Vitronectin antibody (NBP2-20866) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.



Vitronectin antibody detects Vitronectin protein by western blot analysis. Mouse tissue extracts (50 ug) was separated by 10% SDS-PAGE, and the membrane was blotted with Vitronectin antibody (NBP2-20866) diluted by 1:1000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.



Publications

Stuart CH, Riley KR, Boyacioglu O et al. Selection of a Novel Aptamer Against Vitronectin Using Capillary Electrophoresis and Next Generation Sequencing. *Mol Ther Nucleic Acids*. 2016-11-15 [PMID: 27845768] (IF/IHC, Human)

Nystrom A, Thriene K, Mittapalli V et al. Losartan ameliorates dystrophic epidermolysis bullosa and uncovers new disease mechanisms *EMBO Mol Med*. 2015-07-20 [PMID: 26194911] (WB, Human)





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-20866

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-20866

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

