Product Datasheet

WDR62 Antibody NBP2-20897

Unit Size: 0.1 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-20897

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-20897



NBP2-20897

WDR62 Antibody


Unit Size 0.1 ml Concentration Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services. Storage Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles. Clonality Polyclonal Preservative 0.01% Thimerosal Isotype IgG Purity Antigen Affinity-purified Buffer PBS (pH 7), 20% Glycerol, 1% BSA Target Molecular Weight 167 kDa		
Concentration Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services. Storage Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles. Clonality Polyclonal Preservative 0.01% Thimerosal Isotype IgG Purity Antigen Affinity-purified Buffer PBS (pH 7), 20% Glycerol, 1% BSA	Product Information	
contact technical services. Storage Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles. Clonality Polyclonal Preservative 0.01% Thimerosal Isotype IgG Purity Antigen Affinity-purified Buffer PBS (pH 7), 20% Glycerol, 1% BSA	Unit Size	0.1 ml
Clonality Polyclonal Preservative 0.01% Thimerosal Isotype IgG Purity Antigen Affinity-purified Buffer PBS (pH 7), 20% Glycerol, 1% BSA	Concentration	· ·
Preservative 0.01% Thimerosal Isotype IgG Purity Antigen Affinity-purified Buffer PBS (pH 7), 20% Glycerol, 1% BSA	Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
IsotypeIgGPurityAntigen Affinity-purifiedBufferPBS (pH 7), 20% Glycerol, 1% BSA	Clonality	Polyclonal
Purity Antigen Affinity-purified Buffer PBS (pH 7), 20% Glycerol, 1% BSA	Preservative	0.01% Thimerosal
Buffer PBS (pH 7), 20% Glycerol, 1% BSA	Isotype	IgG
- (1),	Purity	Antigen Affinity-purified
Target Molecular Weight 167 kDa	Buffer	PBS (pH 7), 20% Glycerol, 1% BSA
	Target Molecular Weight	167 kDa

Novus Biologicals Rabbit WDR62 Antibody (NBP2-20897) is a polyclonal antibody validated for use in IHC, WB and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.
Rabbit
284403
WDR62
Human, Mouse, Rat
Carrier-protein conjugated synthetic peptide encompassing a sequence within the center region of mouse WDR62. The exact sequence is proprietary.

Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen
Recommended Dilutions	Western Blot 1:500-1:3000, Immunohistochemistry 10 - 1:500, Immunocytochemistry/ Immunofluorescence 1:100-1:1000, Immunohistochemistry-Paraffin 1:100-1:1000, Immunohistochemistry-Frozen 1:100-1:1000


Images

Western Blot: WDR62 Antibody [NBP2-20897] - Whole cell extract (30 ug) was separated by 5% SDS-PAGE, and the membrane was blotted with WDR62 antibody diluted at 1:1000.







Immunocytochemistry/Immunofluorescence: WDR62 Antibody [NBP2-20897] - WDR62 antibody detects WDR62 protein by immunohistochemical analysis. Sample: Frozen-sectioned rat E13.5 brain. Green: WDR62 stained by WDR62 antibody diluted at 1:250. Red: beta Tubulin 3/ Tuj1, stained by beta Tubulin 3/ Tuj1 antibody diluted at 1:500. Blue: Fluoroshield with DAPI.




Immunohistochemistry-Paraffin: WDR62 Antibody [NBP2-20897] - Paraffin-embedded mouse brain diluted at 1:500.




Western Blot: WDR62 Antibody [NBP2-20897] - Sample (30 ug of whole cell lysate) A:NIH-3T3 5% SDS PAGE gel, diluted at 1:1000.




Immunocytochemistry/Immunofluorescence: WDR62 Antibody [NBP2-20897] - WDR62 antibody detects WDR62 protein at nucleus by immunofluorescent analysis. Sample: HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: WDR62 protein stained by WDR62 antibody diluted at 1:5000. Red: phalloidin, a cytoskeleton marker, diluted at 1:100. Scale bar = 10 um.






Immunohistochemistry-Paraffin: WDR62 Antibody [NBP2-20897] - Wdr62 antibody detects Wdr62 protein at nucleus on Ca922 oral cancer xenograft by immunohistochemical analysis. Sample: Paraffin-embedded Ca922 oral cancer xenograft. Wdr62 antibody dilution: 1:500.



Immunohistochemistry-Paraffin: WDR62 Antibody [NBP2-20897] - Paraffin-embedded mouse testis diluted at 1:500.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-20897

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

H00284403-P01-10ug Recombinant Human WDR62 GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-20897

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

