

Product Datasheet

Nanog Antibody (23D2-3C6)

NBP2-22632

Unit Size: 100 ug

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-22632

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-22632



NBP2-22632

Nanog Antibody (23D2-3C6)


Product Information	
Unit Size	100 ug
Concentration	1 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	23D2-3C6
Preservative	0.05% Sodium Azide
Isotype	IgG1
Purity	Protein A purified
Buffer	PBS, 1 mg/mL BSA, 30% glycerol

Product Description	
Host	Mouse
Gene ID	79923
Gene Symbol	NANOG
Species	Human
Marker	Embryonic Stem Cell Marker
Specificity/Sensitivity	It shows specificity to Nanog and is non-reactive to lysates from non-embryonal cell types (e.g. HeLa cell lysate).
Immunogen	Full-length human recombinant protein expressed in bacteria


Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence
Recommended Dilutions	Western Blot 1:500 - 1:2000, Flow Cytometry 1:10 - 1:1000, Immunocytochemistry/ Immunofluorescence 1:50 - 1:100
Application Notes	WB: Detects a prominent approx. 38 kDa protein in embryonal carcinoma cells. Subcellular fractionation shows nuclear localization of Nanog.

Images


Western Blot: Nanog Antibody (23D2-3C6) [NBP2-22632] - Analysis of 60 ug of various whole cell lysates and 10ul of PageRuler Prestained Protein Ladder.




Immunocytochemistry/Immunofluorescence: Nanog Antibody (23D2-3C6) [NBP2-22632] - Analysis of Nanog (green) in H9 embryonic stem cells grown for a few days on Matrigel-coated chamber slides. Cells fixed in 4% paraformaldehyde were permeabilized with 0.1% Triton X-100 for 15 minutes at room temperature. Cells were probed with a Nanog monoclonal antibody at a dilution of 1:200 overnight at 4°C, washed with PBST, and incubated with a fluorescein-conjugated secondary antibody at a dilution of 1:100 for 1 hour at room temperature. Nuclei (blue) were stained with DAPI and cells were analyzed by fluorescence microscopy at 20X magnification.




Flow Cytometry: Nanog Antibody (23D2-3C6) [NBP2-22632] - Analysis of Nanog (blue histogram) on H9 embryonic stem cells. To generate single cells suspensions, colonies were treated with TrypLE cell dissociation enzyme for 5 minutes at 37°C. Cells were incubated with a Nanog monoclonal antibody or mouse IgG (green histogram) at a dilution of 1:100 for 1 hour on ice, washed with PBS + 5% fetal calf serum (FACS buffer), and incubated with a fluorescein-conjugated secondary antibody at a dilution of 1:200 for 30 minutes on ice. Cells were washed with cold FACS buffer, resuspended in 500ul of FACS buffer containing 10ul of 4% paraformaldehyde.




Western Blot: Nanog Antibody (23D2-3C6) [NBP2-22632] - Analysis of 60 ug of NCCIT nuclear and cytoplasmic fractions lysates and 10ul of PageRuler Prestained Protein Ladder




Immunocytochemistry/Immunofluorescence: Nanog Antibody (23D2-3C6) [NBP2-22632] - Analysis of Nanog (green) in NTERA-2 and HeLa cells. Formalin fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 10 minutes at room temperature. Cells were blocked with 1% Blocker BSA for 15 minutes at room temperature. Cells were probed with a Nanog monoclonal antibody at a dilution of 1:50 for at least 1 hour at room temperature, washed with PBS, and incubated with a DyLight 488-conjugated goat anti-mouse IgG secondary antibody. F-Actin (red) was stained with DyLight-554 Phalloidin and nuclei (blue) were stained with Hoechst 33342 dye.



Immunocytochemistry/Immunofluorescence: Nanog Antibody (23D2-3C6) [NBP2-22632] - Analysis of Nanog (green) in HEL 11.4 induced IPS cells grown for a few days on Matrigel-coated chamber slides. Cells fixed in 4% paraformaldehyde were permeabilized with 0.1% Triton X-100 for 15 minutes at room temperature. Cells were probed with a Nanog monoclonal antibody at a dilution of 1:200 overnight at 4C, washed with PBST, and incubated with a fluorescein-conjugated secondary antibody at a dilution of 1:100 for 1 hour at room temperature. Nuclei (blue) were stained with DAPI and cells were analyzed by fluorescence microscopy at 20X magnification.



Flow Cytometry: Nanog Antibody (23D2-3C6) [NBP2-22632] - Analysis of Nanog (blue histogram) on HEL 11.4 induced IPS cells. To generate single cells suspensions, colonies were treated with TrypLE cell dissociation enzyme for 5 minutes at 37C. Cells were incubated with a Nanog monoclonal antibody or mouse IgG (green histogram) at a dilution of 1:100 for 1 hour on ice, washed with PBS + 5% fetal calf serum (FACS buffer), and incubated with a fluorescein-conjugated secondary antibody at a dilution of 1:200 for 30 minutes on ice. Cells were washed with cold FACS buffer, resuspended in 500ul of FACS buffer containing 10ul of 4% paraformaldehyde.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-22632

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)
NBP2-13177PEP	Nanog Antibody Blocking Peptide

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-22632

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

