# **Product Datasheet**

# NPY4R Antibody NLS1071

Unit Size: 0.05 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

**Publications: 1** 

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NLS1071

Updated 4/18/2021 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NLS1071



## **NLS1071**

NPY4R Antibody

Product Information


Product Information	
Unit Size	0.05 ml
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.1% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS
Product Description	
Host	Rabbit
Gene ID	5540
Gene Symbol	NPY4R
Species	Human
Reactivity Notes	Predicted cross-reactivity based on sequence identity: Gibbon (100%), Gorilla (94%), Panda (94%), Monkey (88%), Bovine (88%), Equine (88%), Porcine (88%), Bat (82%), Rabbit (82%).
Specificity/Sensitivity	Human PPYR1. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Immunogen	Synthetic 17 amino acid peptide from C-terminus of human PPYR1.
Product Application Details	
Applications	Immunohistochemistry, Immunohistochemistry-Paraffin

Immunohistochemistry, Immunohistochemistry-Paraffin 4 - 6 ug/ml

# Images


**Recommended Dilutions** 

Immunohistochemistry-Paraffin: NPY4R Antibody [NLS1071] - Anti-NPY4R / PPYR1 antibody IHC of human Lung, Small Cell Carcinoma. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.







Immunohistochemistry-Paraffin: NPY4R Antibody [NLS1071] - Analysis of anti-NPY4R / PPYR1 antibody with small intestine, muscularis propria.




Immunohistochemistry-Paraffin: NPY4R Antibody [NLS1071] - Analysis of anti-NPY4R / PPYR1 antibody with brain at 4 ug/ ml.




Immunohistochemistry-Paraffin: NPY4R Antibody [NLS1071] - Analysis of anti-NPY4R / PPYR1 antibody with human colon, carcinoma.



Immunohistochemistry-Paraffin: NPY4R Antibody [NLS1071] - Anti-NPY4R / PPYR1 antibody IHC of human Skin, Melanoma. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



Immunohistochemistry-Paraffin: NPY4R Antibody [NLS1071] - Anti-PPYR1 antibody IHC of human brain. Immunohistochemistry of formalinfixed, paraffin-embedded tissue after heat-induced antigen retrieval.



## **Publications**

Tarnowski LJ, Kowalec P, Milewski M et al. Nuclear import and export signals of human cohesins SA1/STAG1 and SA2/STAG2 expressed in Saccharomyces cerevisiae. PLoS One 2012-01-01 [PMID: 22715410]



#### **Procedures**

#### Immunohistochemistry protocol for NPY4R Antibody (NLS1071)

Immunohistochemistry Protocol for NPY4R Antibody (NLS1071): Immunohistochemistry

- 1. Prepare tissue with formalin fixation and by embedding it in paraffin wax.
- 2. Make 4-um sections and place on pre-cleaned and charged microscope slides.
- 3. Heat in a tissue-drying oven for 45 minutes at 60 degrees Celcius.
- 4. Deparaffinize the tissues by wash drying the slides in 3 changes of xylene approximately 5 minutes each @ RT.
- 5. Rehydrate the tissues by washing the slides in 3 changes of 100% alcohol approximately 3 minutes each @ RT.
- 6. Wash the slides in 2 changes of 95% alcohol approximately 3 minutes each @ RT.
- 7. Wash the slides in 1 change of 80% alcohol approximately 3 minutes @ RT.
- 8. Rinse the slides in gentle running distilled water approximately 5 minutes @ RT.
- 9. Perform antigen retrieval by steaming the slides in 0.01M sodium citrate buffer (pH 6.0) @ 99-100 degrees Celcius for 20 minutes.
- 10. Remove the slides from the heat and let stand in buffer @ RT for 20 minutes.
- 11. Rinse the slides in 1X TBS-T for 1 minute @ RT.
- \*\*Do not allow the tissues to dry at any time during the staining procedure\*\*
- 12. Begin the immunostaining by applying a universal protein block approximately 20 minutes @ RT.
- 13. Drain protein block from the slides and apply the diluted primary antibody approximately 45 minutes @ RT.
- 14. Rinse the slide in 1X TBS-T approximately 1 minute @ RT.
- 15. Apply a biotinylated anti-rabbit IgG (H+L) secondary approximately 30 minutes @ RT.
- 16. Rinse the slide in 1X TBS-T approximately 1 minute at RT.
- 17. Apply an alkaline phosphatase steptavidin approximately 30 minutes at RT.
- 18. Rinse the slide in 1X TBS-T approximately 1 minute at RT.
- 19. Apply an alkaline phosphatase chromagen substrate approximately 30 minutes at RT.
- 20. Rinse the slide in distilled water approximately 1 minute @ RT.
- \*\*This method should only be used if the chromagen substrate is alcohol insoluble (ie: Vector Red, DAB)\*\*
- 21. Dehydrate the tissue by washing the slides in 2 changes of 80% alcohol approximately 1 minute each @ RT.
- 22. Wash the slides in 2 changes of 95% alcohol approximately 1 minute each @ RT.
- 23. Wash the slides in 3 changes of 100% alcohol approximately 1 minute each @ RT.
- 24. Wash the slides in 3 changes of xyleneapproximately 1 minute each @ RT.
- 25. Apply cover slip.





# **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

# **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

## **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

## **General Contact Information**

www.novusbio.com Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NLS1071

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

