Product Datasheet

TRHR Antibody NLS1200

Unit Size: 0.05 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NLS1200

Updated 6/30/2021 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NLS1200



NLS1200

TRHR Antibody


TRAK AHIIDOOY	
Product Information	
Unit Size	0.05 ml
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.1% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS
Product Description	
Host	Rabbit
Gene ID	7201
Gene Symbol	TRHR
Species	Human, Canine, Equine, Primate, Monkey, Sheep
Reactivity Notes	Gorilla, Gibbon, Marmoset. Predicted cross-reactivity based on sequence identity: Panda (100%), Elephant (94%), Mouse (88%), Bovien (88%), Rabbit (88%), Opossum (88%), Hamster (82%), Turkey (82%), Medaka (82%).
Specificity/Sensitivity	Human TRH Receptor. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Immunogen	Synthetic 17 amino acid peptide from C-terminus of human TRH Receptor.
Product Application Details	
Applications	Immunohistochemistry, Immunohistochemistry-Paraffin

Product Application Details	
Applications	Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry, Immunohistochemistry-Paraffin 1:10 - 1:500




Images


Immunohistochemistry-Paraffin: TRHR Antibody [NLS1200] - Anti-TRH Receptor / TRHR antibody IHC of human Ovary, Carcinoma. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



Immunohistochemistry-Paraffin: TRHR Antibody [NLS1200] - Analysis of anti-TRH Receptor / TRHR antibody with brain, neurons and glia using this antibody at 16 ug/ ml.



Immunohistochemistry-Paraffin: TRHR Antibody [NLS1200] - Analysis of anti-TRH Receptor / TRHR antibody with human lung, non-small cell carcinoma.



Publications

Xiao Y, Segal MR. Biological sequence classification utilizing positive and unlabeled data. Bioinformatics 2008-05-01 [PMID: 18344247]



Procedures

Immunohistochemistry Protocol for Thyrotropin-Releasing Hormone Receptor Antibody (NLS1200)

Immunohistochemistry Protocol for Thyrotropin-Releasing Hormone Receptor Antibody (NLS1200): Immunohistochemistry

- 1. Prepare tissue with formalin fixation and by embedding it in paraffin wax.
- 2. Make 4 um sections and place on pre-cleaned and charged microscope slides.
- 3. Heat in a tissue-drying oven for 45 minutes @ 60 degrees Celcius.
- 4. Deparaffinize the tissues by wash drying the slides in 3 changes of xylene for 5 minutes each @ RT.
- 5. Rehydrate the tissues by washing the slides in 3 changes of 100% alcohol for 3 minutes each @ RT.
- 6. Wash the slides in 2 changes of 95% alcohol for 3 minutes each @ RT.
- 7. Wash the slides in 1 change of 80% alcohol for 3 minutes @ RT.
- 8. Rinse the slides in gentle running distilled water for 5 minutes @ RT.
- 9. Perform antigen retrieval by steaming the slides in 0.01M sodium citrate buffer (pH 6.0) @ 99-100 degrees Celcius for 20 minutes.
- 10. Remove the slides from the heat and let stand in buffer @ RT for 20 minutes.
- 11. Rinse the slides in 1X TBS-T for 1 minute @ RT.
- **Do not allow the tissues to dry at any time during the staining procedure**
- 12. Begin the immunostaining by applying a universal protein block for 20 minutes @ RT.
- 13. Drain protein block from the slides and apply the diluted primary antibody for 45 minutes @ RT.
- 14. Rinse the slide in 1X TBS-T for 1 minute @ RT.
- 15. Apply a biotinylated anti-rabbit IgG (H+L) secondary for 30 minutes @ RT.
- 16. Rinse the slide in 1X TBS-T for 1 minute @ RT.
- 17. Apply an alkaline phosphatase steptavidin for 30 minutes @ RT.
- 18. Rinse the slide in 1X TBS-T for 1 minute @ RT.
- 19. Apply an alkaline phosphatase chromagen substrate for 30 minutes @ RT.
- 20. Rinse the slide in distilled water for 1 minute @ RT.
- **This method should only be used if the chromagen substrate is alcohol insoluble (ie: Vector Red, DAB)**
- 21. Dehydrate the tissue by washing the slides in 2 changes of 80% alcohol for 1 minute each @ RT.
- 22. Wash the slides in 2 changes of 95% alcohol for 1 minute each @ RT.
- 23. Wash the slides in 3 changes of 100% alcohol for 1 minute each @ RT.
- 24. Wash the slides in 3 changes of xylene for 1 minute each @ RT.
- 25. Apply cover slip.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NLS1200

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

