

Product Datasheet

Frizzled-6 Antibody NLS4481

Unit Size: 0.05 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NLS4481

Updated 6/20/2021 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NLS4481



NLS4481

Frizzled-6 Antibody

Product Information

Unit Size	0.05 ml
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.1% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS

Product Description


Host	Rabbit
Gene ID	8323
Gene Symbol	FZD6
Species	Human
Reactivity Notes	Predicted cross-reactivity based on sequence identity: Gorilla (94%), Gibbon (94%), Marmoset (94%), Porcine (88%), Opossum (88%), Bat (82%), Rabbit (82%).
Specificity/Sensitivity	Human Frizzled-6. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except MTMR10 (47%).
Immunogen	Synthetic 17 amino acid peptide from N-terminal extracellular domain of human Frizzled-6.

Product Application Details


Applications	Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry, Immunohistochemistry-Paraffin 14 ug/ml

Images


Immunohistochemistry-Paraffin: Frizzled-6 Antibody [NLS4481] - Analysis of anti-Frizzled 6 / FZD6 antibody with human skeletal muscle at dilution 14 ug/ml.




Immunohistochemistry-Paraffin: Frizzled-6 Antibody [NLS4481] -
Analysis of anti-Frizzled 6 / FZD6 antibody with human prostate.



Immunohistochemistry-Paraffin: Frizzled-6 Antibody [NLS4481] -
Analysis of anti-Frizzled 6 / FZD6 antibody with human skeletal muscle.



Immunohistochemistry-Paraffin: Frizzled-6 Antibody [NLS4481] -
Analysis of anti-Frizzled 6 / FZD6 antibody with human liver at dilution 14
ug/ml.



Immunohistochemistry-Paraffin: Frizzled-6 Antibody [NLS4481] -
Analysis of anti-Frizzled 6 / FZD6 antibody with human pancreas.



Procedures

Immunohistochemistry protocol for Frizzled-6 Antibody (NLS4481)

Immunohistochemistry Protocol for Frizzled-6 Antibody (NLS4481):

Immunohistochemistry

1. Prepare tissue with formalin fixation and by embedding it in paraffin wax.
2. Make 4 um sections and place on pre-cleaned and charged microscope slides.
3. Heat in a tissue-drying oven for 45 minutes @ 60 degrees Celcius.
4. Deparaffinize the tissues by wash drying the slides in 3 changes of xylene for 5 minutes each @ RT.
5. Rehydrate the tissues by washing the slides in 3 changes of 100% alcohol for 3 minutes each @ RT.
6. Wash the slides in 2 changes of 95% alcohol for 3 minutes each @ RT.
7. Wash the slides in 1 change of 80% alcohol for 3 minutes @ RT.
8. Rinse the slides in gentle running distilled water for 5 minutes @ RT.
9. Perform antigen retrieval by steaming the slides in 0.01M sodium citrate buffer (pH 6.0) @ 99-100 degrees Celcius for 20 minutes.
10. Remove the slides from the heat and let stand in buffer @ RT for 20 minutes.
11. Rinse the slides in 1X TBS-T for 1 minute @ RT.

Do not allow the tissues to dry at any time during the staining procedure

12. Begin the immunostaining by applying a universal protein block for 20 minutes @ RT.
13. Drain protein block from the slides and apply the diluted primary antibody for 45 minutes @ RT.
14. Rinse the slide in 1X TBS-T for 1 minute @ RT.
15. Apply a biotinylated anti-rabbit IgG (H+L) secondary for 30 minutes @ RT.
16. Rinse the slide in 1X TBS-T for 1 minute @ RT.
17. Apply an alkaline phosphatase streptavidin for 30 minutes @ RT.
18. Rinse the slide in 1X TBS-T for 1 minute @ RT.
19. Apply an alkaline phosphatase chromagen substrate for 30 minutes @ RT.
20. Rinse the slide in distilled water for 1 minute @ RT.

This method should only be used if the chromagen substrate is alcohol insoluble (ie: Vector Red, DAB)

21. Dehydrate the tissue by washing the slides in 2 changes of 80% alcohol for 1 minute each @ RT.
22. Wash the slides in 2 changes of 95% alcohol for 1 minute each @ RT.
23. Wash the slides in 3 changes of 100% alcohol for 1 minute each @ RT.
24. Wash the slides in 3 changes of xylene for 1 minute each @ RT.
25. Apply cover slip.



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NLS4481

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications