Product Datasheet

5-HT7 Antibody NLS663

Unit Size: 0.05 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NLS663

Updated 6/30/2021 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NLS663



NLS663

5-HT7 Antibody


3-1117 Allibody			
Product Information			
Unit Size	0.05 ml		
Concentration	1.0 mg/ml		
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.		
Clonality	Polyclonal		
Preservative	0.1% Sodium Azide		
Isotype	IgG		
Purity	Immunogen affinity purified		
Buffer	PBS		
Product Description			
Host	Rabbit		
Gene ID	3363		
Gene Symbol	HTR7		
Species	Human		
Specificity/Sensitivity	Human 5HT7 Receptor. BLAST analysis of the peptide immunogen showed no homology with other human proteins.		
Immunogen	Synthetic 16 amino acid peptide from C-terminus of human 5HT7 Receptor.		
Product Application Details			
Applications	Immunohistochemistry, Immunohistochemistry-Paraffin		
D	L 1:4 L :4 L 1:4 L :4 D # # 440 4 500		

•	,	• •	•
Product Application Details			
Applications	Immunohistochemistry,	Immunohistochemistry-Paraffin	
Recommended Dilutions	Immunohistochemistry,	Immunohistochemistry-Paraffin 1:10-1:500	




Images

Immunohistochemistry-Paraffin: 5-HT7 Antibody [NLS663] - Human Brain, Cerebellum: Formalin-Fixed, Paraffin-Embedded (FFPE)



Immunohistochemistry-Paraffin: 5-HT7 Antibody [NLS663] - Analysis of anti-5HT7 Receptor antibody with human brain, neurons and astrocytes.



Procedures

Immunohistochemistry Protocol for 5HT7 Receptor Antibody (NLS663)

Immunohistochemistry Protocol for 5HT7 Receptor Antibody (NLS663): Immunohistochemistry

- 1. Prepare tissue with formalin fixation and by embedding it in paraffin wax.
- 2. Make 4 um sections and place on pre-cleaned and charged microscope slides.
- 3. Heat in a tissue-drying oven for 45 minutes @ 60 degrees Celcius.
- 4. Deparaffinize the tissues by wash drying the slides in 3 changes of xylene for 5 minutes each @ RT.
- 5. Rehydrate the tissues by washing the slides in 3 changes of 100% alcohol for 3 minutes each @ RT.
- 6. Wash the slides in 2 changes of 95% alcohol for 3 minutes each @ RT.
- 7. Wash the slides in 1 change of 80% alcohol for 3 minutes @ RT.
- 8. Rinse the slides in gentle running distilled water for 5 minutes @ RT.
- Perform antigen retrieval by steaming the slides in 0.01M sodium citrate buffer (pH 6.0) @ 99-100 degrees Celcius for 20 minutes.
- 10. Remove the slides from the heat and let stand in buffer @ RT for 20 minutes.
- 11. Rinse the slides in 1X TBS-T for 1 minute @ RT.
- **Do not allow the tissues to dry at any time during the staining procedure**
- 12. Begin the immunostaining by applying a universal protein block for 20 minutes @ RT.
- 13. Drain protein block from the slides and apply the diluted primary antibody for 45 minutes @ RT.
- 14. Rinse the slide in 1X TBS-T for 1 minute @ RT.
- 15. Apply a biotinylated anti-rabbit IgG (H+L) secondary for 30 minutes @ RT.
- 16. Rinse the slide in 1X TBS-T for 1 minute @ RT.
- 17. Apply an alkaline phosphatase steptavidin for 30 minutes @ RT.
- 18. Rinse the slide in 1X TBS-T for 1 minute @ RT.
- 19. Apply an alkaline phosphatase chromagen substrate for 30 minutes @ RT.
- 20. Rinse the slide in distilled water for 1 minute @ RT.
- **This method should only be used if the chromagen substrate is alcohol insoluble (ie: Vector Red, DAB)**
- 21. Dehydrate the tissue by washing the slides in 2 changes of 80% alcohol for 1 minute each @ RT.
- 22. Wash the slides in 2 changes of 95% alcohol for 1 minute each @ RT.
- 23. Wash the slides in 3 changes of 100% alcohol for 1 minute each @ RT.
- 24. Wash the slides in 3 changes of xylene for 1 minute each @ RT.
- 25. Apply cover slip.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada 21 Canmotor Ave

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NLS663

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

