

Product Datasheet

CXCR4 Antibody - BSA Free

NB100-56437SS

Unit Size: 0.025 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 1 Publications: 21

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-56437

Updated 2/21/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-56437



NB100-56437SS


CXCR4 Antibody - BSA Free

Product Information	
Unit Size	0.025 mg
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein G purified
Buffer	PBS
Product Description	
Host	Rabbit
Gene ID	7852
Gene Symbol	CXCR4
Species	Human, Mouse, Rat, Sheep
Immunogen	Rabbit anti-CXCR4 polyclonal antibody was raised against a peptide corresponding to amino acids 328-338 of human CXCR4.
Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin, Knockdown Validated
Recommended Dilutions	Western Blot 1 - 2 ug/mL, Immunohistochemistry 1:10 - 1:500, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 1:10 - 1:500. Use reported in scientific literature (Zagzag (2005)), Immunohistochemistry-Frozen reported in scientific literature (Nimmagadda et al (2010)), Knockdown Validated
Application Notes	In HeLa, a 45 kDa band is observed.




Images


Western Blot: CXCR4 Antibody [NB100-56437] - Western blot shows lysates of Jurkat human T lymphocyte parental cell line and CXCR4 knockout (KO) Jurkat cell line. PVDF membrane was probed with 1 ug/ml of Rabbit Anti-Human CXCR4 Polyclonal Antibody (Catalog # NB100-56437) followed by HRP-conjugated Anti-Rabbit IgG Secondary Antibody (Catalog #HAF008). Specific band was detected for CXCR4 at approximately 38 kDa (as indicated) in the parental Jurkat cell line, but is not detectable in the knockout Jurkat cell line. This experiment was conducted under reducing conditions.




Western Blot: CXCR4 Antibody [NB100-56437] - Detection of 15 ug in HeLa cells with antibody concentration at 1 ug/mL.



Immunocytochemistry/Immunofluorescence: CXCR4 Antibody [NB100-56437] - Jurkat cells were fixed for 10 minutes using 10% formalin and then permeabilized for 5 minutes using 1X TBS + 0.5% Triton X-100. The cells were incubated with anti-CXCR4 (NB100-56437) at a 1:25 dilution for 1 hour at room temperature and detected with an anti-rabbit DyLight 488 (Green) at a 1:500 dilution. Alpha tubulin was used as a co-stain at a 1:1000 dilution and detected with and anti-mouse DyLight 550 (Red) at a 1:500 dilution. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 40X objective.



Immunohistochemistry-Paraffin: CXCR4 Antibody [NB100-56437] - CXCR4 was detected in immersion sections of human tonsil using rabbit anti-human antibody (Catalog # NB100-56437) at 1:300 dilution overnight at 4C. Tissue was stained using the VisuCyte anti-rabbit HRP polymer detection reagent with DAB chromogen (brown) and counterstained with hematoxylin (blue).



Publications

Hiraide T, Tsuda N, Momoi M et al. CXCL12/CXCR4 pathway as a novel therapeutic target for RNF213 -associated pulmonary arterial hypertension Scientific Reports 2024-11-04 [PMID: 39496725]

Lyu C, Ye Y, Lensing MM et al. Targeting Gi/o protein-coupled receptor signaling blocks HER2-induced breast cancer development and enhances HER2-targeted therapy JCI Insight 2021-09-22 [PMID: 34343132] (Immunohistochemistry)

Locri F, Dal Monte M, Aronsson M et al. UPARANT is an effective antiangiogenic agent in a mouse model of rubeosis iridis Journal of Molecular Medicine 2019-09-01 [PMID: 31243519] (In vivo assay)

Queen NJ, Deng H, Huang W et al. Environmental Enrichment Mitigates Age-Related Metabolic Decline and Lewis Lung Carcinoma Growth in Aged Female Mice Cancer Prevention Research 2021-12-01 [PMID: 34535449] (Western Blot)

Yamamoto Y, Kadoya K, Terkawi MA et al. Neutrophils delay repair process in Wallerian degeneration by releasing NETs outside the parenchyma Life science alliance 2022-10-01 [PMID: 35961782] (IHC-P, Mouse)

Zheng M, Oh SH, Choi N et al. CXCL12 inhibits hair growth through CXCR4 Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie 2022-04-21 [PMID: 35462338] (IF/IHC, Mouse)

Irie Y, Tsubota M, Ishikura H et al. Macrophage-derived HMGB1 as a Pain Mediator in the Early Stage of Acute Pancreatitis in Mice: Targeting RAGE and CXCL12/CXCR4 Axis J Neuroimmune Pharmacol 2017-07-28 [PMID: 28755135] (WB, Mouse)

Pelagalli A, Nardelli A, Lucarelli E et al. Autocrine signals increase Ovine Mesenchymal Stem Cells migration through Aquaporin-1 and CXCR4 overexpression J. Cell. Physiol. 2018-01-18 [PMID: 29345324] (WB)

Gagner JP, Sarfraz Y, Ortenzi V et al. Multifaceted C-X-C Chemokine Receptor 4 Inhibition Interferes with Anti-Vascular Endothelial Growth Factor Therapy-Induced Glioma Dissemination. Am. J. Pathol. 2017-07-20 [PMID: 28734730] (WB, Mouse)

Hogg K, Etherington SL, Young JM et al. Inhibitor of differentiation (Id) genes are expressed in the steroidogenic cells of the ovine ovary and are differentially regulated by members of the transforming growth factor-beta family. Endocrinology [PMID: 20032053]

Details:

Used the Azide Free form of this antibody.

Dieckow J, Brandt W, Hattermann K et al. CXCR4 and CXCR7 Mediate TFF3-Induced Cell Migration Independently From the ERK1/2 Signaling Pathway. Invest. Ophthalmol. Vis. Sci. 2016-01-01 [PMID: 26780310] (ICC/IF, Human)

Nimmagadda S, Pullambhatla M, Stone K et al. Molecular imaging of CXCR4 receptor expression in human cancer xenografts with [64Cu]AMD3100 positron emission tomography. Cancer Res. 2010-05-15 [PMID: 20460522]

More publications at <http://www.novusbio.com/NB100-56437>





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-56437

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

