

Product Datasheet

IKK beta Antibody (10A9B6) - BSA Free NB100-56513SS

Unit Size: 0.025 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 1 Publications: 9

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-56513

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-56513



NB100-56513SS


IKK beta Antibody (10A9B6) - BSA Free

Product Information	
Unit Size	0.025 mg
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	10A9B6
Preservative	0.05% Sodium Azide
Isotype	IgG1 Kappa
Purity	Protein G purified
Buffer	PBS
Product Description	
Description	Novus Biologicals Mouse IKK beta Antibody (10A9B6) - BSA Free (NB100-56513) is a monoclonal antibody validated for use in IHC, WB, Flow, ICC/IF, Simple Western and IP. Anti-IKK beta Antibody: Cited in 9 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	3551
Gene Symbol	IKBKB
Species	Human, Mouse
Immunogen	Full-length human IKKb/IKK2 recombinant protein was used as immunogen (NP_001547).
Product Application Details	
Applications	Western Blot, Simple Western, Flow Cytometry, Flow (Intracellular), Gel Super Shift Assays, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunoprecipitation
Recommended Dilutions	Western Blot 2 - 5 ug/mL, Simple Western 1:12.5, Flow Cytometry 1:10 - 1:1000, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence reported in scientific literature (PMID 24825920), Immunoprecipitation 2 - 5 ug / 10 ⁶ cells, Immunohistochemistry-Frozen reported by customer review, Gel Super Shift Assays reported in scientific literature (PMID 19662631), Flow (Intracellular)
Application Notes	An approx. 87 kDa band should be observed. In Simple Western only 10 - 15 uL of the recommended dilution is used per data point. See Simple Western Antibody Database for Simple Western validation: Tested in HeLa lysate 1.0 mg/mL, separated by Size, antibody dilution of 1:12.5, apparent MW was 89 kDa. Separated by Size-Wes, Sally Sue/Peggy Sue.




Images


Western Blot: IKK beta Antibody (10A9B6) [NB100-56513] - IKK beta in Daudi cell lysate using IKK beta antibody at 5 ug/ml.




Immunohistochemistry-Frozen: IKK beta Antibody (10A9B6) [NB100-56513] - Mouse skin tissue section stained with IKK beta antibody. IHC-Fr image submitted by a verified customer review.




Flow (Intracellular): IKK beta Antibody (10A9B6) [NB100-56513] - Analysis using the Alexa Fluor (R) 488 conjugate. Staining of IKK in HEK 293 cells using 0.1 ug of this antibody. Green histogram represents the isotype control, red represents the IKK antibody. Image using the Azide Free format of this antibody.



Simple Western: IKK beta Antibody (10A9B6) [NB100-56513] - Analysis using Azide Free version of NB100-56513. Lane view shows a specific band for IKK beta in 1.0 mg/ml of HeLa lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system. *Non-specific interaction with the



Immunoprecipitation: IKK beta Antibody (10A9B6) [NB100-56513] - Antibody was used to immunoprecipitate IKKb from 1×10^6 Daudi cells and the immuno-precipitated protein was detected by western blotting using NB100-56509. IKKb was detected as a protein of ~87 kD. Image using the Azide Free format of this antibody.



Publications

Yang Gong, Rosen Daniel G, Liu Guangzhi et al. CXCR2 promotes ovarian cancer growth through dysregulated cell cycle, diminished apoptosis, and enhanced angiogenesis. *Clin Cancer Res.* 2010-08-01 [PMID: 20505188] (WB)

Monsalve Eva, Ruiz-Garcia Almudena, Baladron Victoriano et al. Notch1 upregulates LPS-induced macrophage activation by increasing NF-kappaB activity. *Eur J Immunol.* 2009-09-01 [PMID: 19662631] (EMSA)

Vacca Alessandra, Felli Maria Pia, Palermo Rocco et al. Notch3 and pre-TCR interaction unveils distinct NF-kappaB pathways in T-cell development and leukemia. *EMBO J.* 2006-03-08 [PMID: 16498412] (Mouse)

Tare Nadine, Li Hongli, Morschauser Andrew et al. KU812 cells provide a novel in vitro model of the human IL-33/ST2L axis: functional responses and identification of signaling pathways. *Exp Cell Res.* 2010-09-10 [PMID: 20406635] (WB, Human)

Paul S, Traver MK, Kashyap AK et al. T Cell Receptor Signals to NF-kB Are Transmitted by a Cytosolic p62-Bcl10-Malt1-Ikk Signalosome. *Sci Signal.* 2014-05-14 [PMID: 24825920] (ICC/IF, Mouse)

Details:

D10 mouse T-helper cells, Fig S1B

Lamothe B, Campos AD, Webster WK et al. The RING domain and first zinc finger of TRAF6 coordinate signaling by interleukin-1, lipopolysaccharide, and RANKL. *J Biol Chem.* 2008-09-05 [PMID: 18617513]

Fitzgerald KA, McWhirter SM, Faia KL et al. IKKepsilon and TBK1 are essential components of the IRF3 signaling pathway. *Nat Immunol.* 2003-05-01 [PMID: 12692549] (WB)

Bakkar N, Ladner K, Canan BD et al. IKKa and alternative NF-kB regulate PGC-1B to promote oxidative muscle metabolism. *J Cell Biol.* 2012-02-20 [PMID: 22351927]

Scott MJ, Billiar TR. Beta2-integrin-induced p38 MAPK activation is a key mediator in the CD14/TLR4/MD2-dependent uptake of lipopolysaccharide by hepatocytes. *J Biol Chem.* 2008-10-24 [PMID: 18701460]

Details:

1. MyD88 (IMG-2005): MyD88 peptide inhibition (mouse WT hepatocytes), Figs. 4A,D. 2. TIRAP (IMG-2006): TIRAP peptide inhibition (mouse WT hepatocytes), Figs. 4B,D; 5B; 6E.



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-
techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-56513

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

