### **Product Datasheet**

# Lactate Dehydrogenase Antibody [Biotin] NB600-862-0.1mg

Unit Size: 0.1 mg

Store lyophilized antibody at 4C in the dark. Aliquot reconstituted liquid and store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB600-862

Updated 10/23/2024 v.20.1

## Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NB600-862



#### NB600-862-0.1mg

Lactate Dehydrogenase Antibody [Biotin]

Lactate Dehydrogenase Antibody [Biotin]	
Product Information	
0.1 mg	
LYOPH mg/ml	
Store lyophilized antibody at 4C in the dark. Aliquot reconstituted liquid and store at -20C. Avoid freeze-thaw cycles.	
Polyclonal	
0.01% Sodium Azide	
Reconstitute with 100 ul deionized water (or equivalent)	
IgG	
Biotin	
Multi-step	
Lyophilized from 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free	
Product Description	
Store vial at 4C prior to restoration. For extended storage aliquot contents and freeze at -20C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4C as an undiluted liquid. Dilute only prior to immediate use.  This antibody is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Goat Serum as well as purified and partially purified Lactate Dehydrogenase [Rabbit Muscle]	
Goat	
3945	
LDHB	
Human, Rabbit	
BLAST analysis was used to determine that cross reactivity is suggested for both muscle and heart isoforms (LDH-A and LDH-B) from most mammalian species	
BLAST analysis was used to determine that cross reactivity is suggested for both muscle and heart isoforms (LDH-A and LDH-B) from most mammalian species.	
This Lactate Dehydrogenase Antibody was prepared from whole goat serum produced by repeated immunizations with a full length Lactate Dehydrogenase protein isolated from rabbit muscle. (Uniprot: P13491)	
Product Application Details	
Western Blot, Dot Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation	
Western Blot 1:500-1:5000, ELISA 1:5000-1:20000, Immunohistochemistry 1:10-1:500, Immunocytochemistry/ Immunofluorescence 1:10-1:2000, Immunoprecipitation 1:100, Immunohistochemistry-Paraffin 1:2500-1:5000, Dot Blot	




#### **Application Notes**


This product has been tested by ELISA, Western blot, and immunofluorescence. This antibody is assayed against 1.0 ug of Lactate Dehydrogenase in a standard capture ELISA using Peroxidase Conjugated Streptavidin and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:20,000 of the reconstitution concentration is suggested for this product.

#### **Images**


Western Blot: Lactate Dehydrogenase Antibody [Biotin] [NB600-862] - Lane 1-4: HeLa cell extracts cytoplasmic fraction (CF). Lane 5-8: HeLa cell extracts nuclear fraction (NF). Load: 30 ug per lane. Primary antibody: LDH antibody at 1:400 for overnight at 4C. Secondary antibody: IRDye800 secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO/TBST overnight at 4C. Predicted/Observed size: 36.6 kDa, 36 kDa for LDH. Other band(s): None.



Immunocytochemistry/Immunofluorescence: Lactate Dehydrogenase Antibody [Biotin] [NB600-862] - Tissue: HeLa cells. Fixation: fixed for 5 min in 1:1 MeTOH:Acetone, blocked with incubated with blocking buffer for 15 min. Antigen retrieval: not required. Primary antibody: Lactate Dehydrogenase antibody at 1:200 for 1 h at RT. Secondary antibody: DyLight 488 conjugated Streptavidin antibody at 1:10,000 for 30 min at RT. Staining: Lactate Dehydrogenase as green fluorescent signal.



Western Blot: Lactate Dehydrogenase Antibody [Biotin] [NB600-862] - Lane 1: Lactate Dehydrogenase (Rabbit Muscle). Load: 50 ng per lane. Primary antibody: Goat anti-Lactate Dehydrogenase Antibody Biotin Conjugated at 1:1,000 overnight at 4C. Secondary antibody: HRP streptavidin secondary antibody at 1:40,000 for 30 min at RT. Block: incubated with blocking buffer for 30 min at RT. Predicted/Observed size: 36.5 kDa, 36.5 kDa for Lactate Dehydrogenase.





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

#### Products Related to NB600-862-0.1mg

NBP2-29370 Streptavidin Native Protein

BAF108 Goat IgG Isotype Control [Biotin]

NBP3-24510 Lactate Dehydrogenase Activity Assay Kit (Colorimetric)

NB300-221 GAPDH Antibody (1D4)

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB600-862

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

