

Product Datasheet

Centaurin beta 2 Antibody

NBP2-24528

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-24528

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-24528



NBP2-24528**Centaurin beta 2 Antibody**

Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS and 0.05% BSA


Product Description	
Description	Novus Biologicals Rabbit Centaurin beta 2 Antibody (NBP2-24528) is a polyclonal antibody validated for use in IHC and WB. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	23527
Gene Symbol	ACAP2
Species	Human, Mouse, Rat, Canine, Equine, Primate
Reactivity Notes	Immunogen displays the following percentage of sequence identity for non-tested species: Monkey (87%).
Immunogen	A portion of amino acids 350-400 of human ACAP2 was used as the immunogen for the antibody.

Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry
Recommended Dilutions	Western Blot 0.25-0.5 ug/ml, Immunohistochemistry, Immunohistochemistry-Paraffin 10 ug/ml




Images

Western Blot: Centaurin beta 2 Antibody [NBP2-24528] - analysis of ACAP2 in human testis lysate in the 1) absence and 2) presence of immunizing peptide, 3) mouse testis lysate and 4) rat testis lysate using ACAP2 antibody at 0.25 ug/ml. I goat anti-rabbit Ig HRP secondary antibody and PicoTest ECL substrate solution were used for this test.



Immunohistochemistry-Paraffin: Centaurin beta 2 Antibody [NBP2-24528] - Formalin-fixed, paraffin-embedded human prostate stained with ACAP2 antibody at 10 ug/ml. Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM sodium citrate buffer, pH 6.0 for 10-20 min followed by cooling at RT for 20 min.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-24528

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
H00023527-P01-10ug	Recombinant Human Centaurin beta 2 GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-24528

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

