

Product Datasheet

TLR8 Antibody (44C143) - BSA Free NBP2-24917SS

Unit Size: 0.025 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 52

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-24917

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-24917



NBP2-24917SS


TLR8 Antibody (44C143) - BSA Free

Product Information	
Unit Size	0.025 mg
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	44C143
Preservative	0.05% Sodium Azide
Isotype	IgG1 Kappa
Purity	Protein G purified
Buffer	PBS
Product Description	
Description	Novus Biologicals Mouse TLR8 Antibody (44C143) - BSA Free (NBP2-24917) is a monoclonal antibody validated for use in IHC, WB, Flow, ICC/IF and Simple Western. Anti-TLR8 Antibody: Cited in 49 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	51311
Gene Symbol	TLR8
Species	Human, Mouse, Rabbit
Immunogen	This antibody was developed against a KLH-conjugated synthetic peptide of human TLR8, within amino acids 750-850.
Product Application Details	
Applications	Western Blot, Simple Western, Immunohistochemistry-Paraffin, Dot Blot, Flow Cytometry, Flow (Cell Surface), Flow (Intracellular), Immunocytochemistry/Immunofluorescence, Immunohistochemistry, CyTOF-ready
Recommended Dilutions	Western Blot 1-3 ug/ml, Simple Western 20 ug/mL, Flow Cytometry 0.5-1 ug/10 ⁶ cells, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin 5 ug/ml, Dot Blot reported in scientific literature (PMID 27248820), Flow (Cell Surface) reported in multiple pieces of scientific literature, Flow (Intracellular) reported in reported in multiple pieces of scientific literature (PMID 24836676), CyTOF-ready
Application Notes	Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM sodium citrate buffer, pH 6.0 for 10-20 min followed by cooling at RT for 20 min. This antibody is CyTOF ready. See Simple Western Antibody Database for Simple Western validation: Tested in Ramos and Human Spleen lysate 0.5 mg/mL, separated by Size, antibody dilution of 20 ug/mL




Images


Flow Cytometry: TLR8 Antibody (44C143) [NBP2-24917] - Intracellular expression of TLR8 in monocyte-derived macrophages (MDMs). MDMs which were incubated with PE-conjugated anti-TLR8 or isotype antibody control. Intracellular expression was determined by flow cytometry. Representative profiles were similar in four independent experiments (n=4 subjects). Image collected and cropped by CiteAb from the following publication ([//doi.org/10.1371/journal.pone.0104039](https://doi.org/10.1371/journal.pone.0104039)) licensed under a CC-BY license.




Flow Cytometry: TLR8 Antibody (44C143) [NBP2-24917] - Expression of TLR8 in Fravel, L363, OPM1, RPMI8226, XG1, and NCI-H929 as determined by flow cytometry. HCMLs were stained using an intracellular staining protocol with TLR8 antibodies followed by relevant secondary fluorescent-conjugated antibodies. Filled histograms (purple) represent the isotype controls and the open histograms (red) indicate TLR8. Data are representative for analysis of ≥ 2 independent experiments. Image collected and cropped by CiteAb from the following publication ([//doi.org/10.1371/journal.pone.0060671](https://doi.org/10.1371/journal.pone.0060671)) licensed under a CC-BY license.




Western Blot: TLR8 Antibody (44C143) [NBP2-24917] - All the cell lines expressed mRNA for TLR8 (Figure 1) and displayed a strong expression of TLR8 protein (Figure 2). Lysates of human intestinal tissue and mouse splenocytes were used as positive controls. Beta-actin was served as loading control and was used to normalize expression levels between cells. Data are representative for analysis of ≥ 2 independent experiments. Image collected and cropped by CiteAb from the following publication ([//doi.org/10.1371/journal.pone.0060671](https://doi.org/10.1371/journal.pone.0060671)) licensed under a CC-BY license.




Immunohistochemistry-Paraffin: TLR8 Antibody (44C143) [NBP2-24917] - Formalin-fixed, paraffin-embedded human spleen stained with TLR8 antibody at 2 ug/ml.




Flow Cytometry: TLR8 Antibody (44C143) [NBP2-24917] - HEK cells expressing TLR8 were treated with ssRNA40-Alexa 546 followed by fixation, permeabilization and staining for TLR8 using anti-TLR8 Alexa 488 conjugate or isotype control IgG1/Kappa Alexa 488. TLR8 detection is shown along the horizontal axis, and ssRNA40 Alexa 546 detection is shown along the vertical axis. Direct excitation (488 and 561 nm lasers) of both fluorophores (Alexa 488 and Alexa 546) allowed quantitation of cells staining positive for TLR8 (anti-TLR8-Alexa488) and cells that had taken up ssRNA40-Alexa 546. Single-stranded RNA40 binds TLR8 (flow cytometry FRET assay). A representative experiment is shown from three independent experiments with similar results. Image collected and cropped by CiteAb from the following publication ([//doi.org/10.1371/journal.pone.0104039](https://doi.org/10.1371/journal.pone.0104039)) licensed under a CC-BY license.



Western Blot: TLR8 Antibody (44C143) [NBP2-24917] - Analysis of TLR8 in human Ramos cell lysate. Goat anti-mouse Ig HRP secondary antibody and PicoTect ECL substrate solution were used for this test.



Simple Western: TLR8 Antibody (44C143) [NBP2-24917] - Simple Western lane view shows a specific band for TLR8 in 0.5 mg/ml of Ramos (left) and Human Spleen (right) lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system. * Non-specific interaction with the 230 kDa Simple Western standard may be seen with this antibody. Image from the standard format of this antibody.



Publications

Gowda M, Venkatraman A, Mechler-Hickson A et al. Toll Like Receptors in the Larynx: Implications for Laryngeal Immunology. The Laryngoscope 2025-06-23 [PMID: 40546085]

Kumbol V, Ivanov A, McGurran H et al. Neurodegenerative disease-associated microRNAs acting as signaling molecules modulate CNS neuron structure and viability. Cell communication and signaling : CCS 2025-04-25 [PMID: 40275260]

Hiltunen N, Kemi N, Väyrynen JP et Al. Toll-like receptors 1-9 in small bowel neuroendocrine tumors-Clinical significance and prognosis PLoS One 2024-05-06 [PMID: 38709790]

Grant R. Campbell, Pratima Rawat, Stephen A. Spector Pacritinib Inhibition of IRAK1 Blocks Aberrant TLR8 Signalling by SARS-CoV-2 and HIV-1-Derived RNA Journal of Innate Immunity 2023-01-01 [PMID: 35785771]

Campbell GR, Rawat P, Teodorof-Diedrich C, Spector SA IRAK1 inhibition blocks the HIV-1 RNA mediated pro-inflammatory cytokine response from microglia The Journal of general virology 2023-05-01 [PMID: 37256770]

Helminen O, Huhta H, Lehenkari Petri P. Nucleic acid-sensing Toll-like receptors 3, 7 and 8 in esophageal epithelium, Barrett's esophagus, dysplasia and adenocarcinoma Oncoimmunology 2016-07-29 [PMID: 27467941]

de Barros De Luccia T, Ono E, Menon R et al. The effect of Gestational Diabetes Mellitus on the fetal compartment Journal of Reproductive Immunology 2021-03-01 [PMID: 33836321] (IHC-P, Human)

Li Y, Chen X, Jin R et al. Injectable hydrogel with MSNs/microRNA-21-5p delivery enables both immunomodification and enhanced angiogenesis for myocardial infarction therapy in pigs Science advances 2021-02-05 [PMID: 33627421] (WB, Mouse)

Chatillon JF Direct Toll-Like Receptor 8 signaling increases the functional avidity of human CD8+ T lymphocytes generated for adoptive T cell therapy strategies Immun Inflamm Dis. 2015-03-01 [PMID: 25866635] (FLOW, Human)

Details:

Citation using the PE format of this antibody.

Menendez D, Snipe J, Marzec J et al. p53-responsive TLR8 SNP enhances human innate immune response to respiratory syncytial virus J. Clin. Invest. 2019-08-20 [PMID: 31430261] (FLOW)

Evankovich J, Lear T, Baldwin C et al. Toll-Like Receptor 8 Stability is Regulated by Ring Finger 216 in response to circulating MicroRNAs Am. J. Respir. Cell Mol. Biol. 2019-08-06 [PMID: 31385713]

Huhta H, Helminen O, Kauppila JH et al. The Expression of Toll-like Receptors in Normal Human and Murine Gastrointestinal Organs and the Effect of Microbiome and Cancer J. Histochem. Cytochem. 2016-08-01 [PMID: 27370795] (IHC-P, Human)

More publications at <http://www.novusbio.com/NBP2-24917>





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-
techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-24917

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

