

Product Datasheet

Pam3CSK4, TLR1 and TLR2 Ligand NBP2-25297-0.025mg

Unit Size: 0.025 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 1 Publications: 30

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-25297

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-25297



NBP2-25297-0.025mg


Pam3CSK4, TLR1 and TLR2 Ligand

Product Information	
Unit Size	0.025 mg
Concentration	Please see the protocols for proper use of this product. If no protocol is available, contact technical services for assistance.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Buffer	Sterile Water
Product Description	
Description	Pam3CSK4, TLR1 and TLR2 Ligand is a synthetic tripalmitoylated lipoheptapeptide analog of the immunologically active N-terminal portion of bacterial lipoprotein. It activates monocytes and macrophages and is also a potent activator of proinflammatory transcription factor NF- κ B. Pam3CSK4, TLR1 and TLR2 Ligand is recognized by a heterodimer formed between TLR1 and TLR2. Pam3CSK4, TLR1 and TLR2 Ligand stimulation: If your cell line does not naturally express TLR1+ TLR2, co-transfect with plasmids pCMV/TLR1 and pCMV/TLR2. Forty-eight hrs after transfection, stimulate cells with 10 to 100 ng/ml of Pam3CSK4 for 6-24 hrs. Determine Pam3CSK4 stimulation using appropriate detection assays.
Species	Human, Mouse, Bacteria, Chicken
Reactivity Notes	Human reactivity reported in scientific literature (PMID: 24105263). Mouse reactivity reported in scientific literature (PMID: 25511699) Use in Chicken reported in scientific publication (PMID: 32733155). Use in Bacteria reported in scientific publication (PMID: 31482074).
Specificity/Sensitivity	Pam3CSK4, TLR1 and TLR2 Ligand
Product Application Details	
Applications	Functional, In vitro assay, In vivo assay, Ligand Activation
Recommended Dilutions	Functional, In vitro assay, In vivo assay, Ligand Activation




Images


Pam3CSK4, TLR1 and TLR2 Ligand [NBP2-25297] - 293 cells were transfected with pCMV/TLR1-2 plasmid and pNF-kB/SEAP plasmid using Lipofectamin 2000. After 48 hrs of transfection, 50 ng/ml of Pam3CSK4 was added. Cells were incubated at 37C for 24 hrs. Transfected cell supernatant was collected and analyzed using the NF-kB SEAPorter Assay kit. pCMV/SEAP plasmid was used to check transfection efficiency.



Pam3CSK4, TLR1 and TLR2 Ligand [NBP2-25297] - Evaluation of the ligand activity on TLR2/NF-kB SEAPorter HEK 293 cell line. IML-102 cell line is a stably co-transfected cell line that expresses full-length human Toll-like receptor 2 (TLR2) and the secreted alkaline phosphatase reporter gene under the transcriptional control of an NF-KB response element. IML-102 cells were plated in 96-well plates at 5×10^4 cells/well. After 16 h, cells were stimulated with various amounts of for 24 h. SEAP was analyzed using the SEAPorter Assay Kit . Data Summary: specifically activated the TLR2/1-depedent NF-KB/SEAP reporter cells in a dose dependent manner (Note: HEK 293 cells endogenously express TLR1).



Pam3CSK4, TLR1 and TLR2 Ligand [NBP2-25297] - Differentiated human macrophages were treated with or without 50 ng/mL Pam3CSK4 for 2 h, and the release of IL-6 in culture supernatants of cells was measured by ELISA kit.



Publications

Liu Y, Diamond SL. Activation of Most Toll-Like Receptors in Whole Human Blood Attenuates Platelet Deposition on Collagen under Flow Journal of Immunology Research 2023-01-17 [PMID: 36703865] (Block/Neutralize)

Yamamoto M, Miyoshi M, Morioka K et al. Anti-nucleolin aptamer, iSN04, inhibits the inflammatory responses in C2C12 myoblasts by modulating the β -catenin/NF- κ B signaling pathway Biochemical and biophysical research communications 2023-04-26 [PMID: 37127012]

Horii T, Orikawa Y, Ohira Y et al. Peptidoglycan-Like Components in Z-100, Extracted from Mycobacterium tuberculosis Strain Aoyama B, Increase IL-12p40 via NOD2 Journal of Immunology Research 2022-06-22 [PMID: 35785036] (LA)

Kawahara T, Ito A, Kiso A, Kawamoto F Inhibitory effect of strawberry geranium (Saxifraga stolonifera) on Toll-like receptor 2-mediated inflammatory response in human skin keratinocytes Journal of Ethnopharmacology 2021-04-01 [PMID: 33819504]

Terada T, Nii T, Isobe N, Yoshimura Y Effects of Toll-like Receptor Ligands on the Expression of Proinflammatory Cytokines and Avian beta-defensins in Cultured Chick Intestine J Poult Sci 2020-07-25 [PMID: 32733155] (Chicken)

Kanoh H, Nitta T, Go S et al. Homeostatic and pathogenic roles of GM3 ganglioside molecular species in TLR4 signaling in obesity EMBO J. 2020-05-07 [PMID: 32378734] (Mouse)

Effects of TLR Ligands on the Expression of Cytokines and Possible Role of NF κ B in its Process in the Theca of Chicken Follicles Kang Y, Nii T, Isobe N J Poult Sci [PMID: 32055188] (Func, Chicken)

Effects of delayed hypothermia on time-dependent microglial production of inflammatory and anti-inflammatory factors. Matsui T, Kida H, Iha T et al. Folia Neuropathol [PMID: 25118900] (Func, Mouse)

Croasdell A, Sime PJ, Phipps RP Resolvin D2 decreases TLR4 expression to mediate resolution in human monocytes FASEB J. 2016-09-30 [PMID: 27256622] (LA, LA, Human)

Yamaguchi M, Hirose Y, Takemura M, et al Streptococcus pneumoniae Evades Host Cell Phagocytosis and Limits Host Mortality Through Its Cell Wall Anchoring Protein PfbA. Front Cell Infect Microbiol. 2019-08-20 [PMID: 31482074] (Bacteria)

Details:

Mice were infected with *S. pneumoniae*.

Sun L, Hult EM, Cornell TT et al. Loss of myeloid-specific protein phosphatase 2A enhances lung injury and fibrosis and results in IL-10 dependent sensitization of epithelial cell apoptosis Am. J. Physiol. Lung Cell Mol. Physiol. 2019-03-06 [PMID: 30838865] (Func, Mouse)

Nihashi Y, Ono T, Kagami H, Takaya T. Toll-like receptor ligand-dependent inflammatory responses in chick skeletal muscle myoblasts. J. Orthop. Res. 2018-10-31 [PMID: 30389519]

More publications at <http://www.novusbio.com/NBP2-25297>





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Support products are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-25297

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

