

Product Datasheet

Thyroid Peroxidase Antibody

NBP2-27544

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-27544

Updated 5/6/2020 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-27544



NBP2-27544

Thyroid Peroxidase Antibody

Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA

Product Description	
Host	Goat
Gene ID	7173
Gene Symbol	TPO
Species	Human
Specificity/Sensitivity	This antibody is expected to recognize all reported isoforms (NP_000538.3; NP_783650.1; NP_783652.1; NP_783653.1).
Immunogen	Peptide with sequence ELRCGKHQAVGTSPQ corresponding to C-Terminus according to NP_000538.3, NP_783650.1, NP_783652.1, NP_783653.1.

Product Application Details	
Applications	Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Peptide ELISA
Recommended Dilutions	Western Blot 0.1 - 0.3 ug/ml, Immunohistochemistry 5 - 10 ug/ml, Immunohistochemistry-Paraffin 5-10 ug/ml, Peptide ELISA Detection limit 1:32000
Application Notes	WB: Approx. 110 kDa band observed in human thyroid gland lysates (calculated MW of 103 kDa band according to NP_000538.3). IHC-P: Human thyroid gland shows textured staining in the cytoplasm of epithelial cells.

Images

Western Blot: Thyroid Peroxidase Antibody [NBP2-27544] - (0.1 ug/ml) Analysis of Thyroid Gland lysate (35 ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

250kDa
150kDa
100kDa
75kDa
50kDa
37kDa
25kDa
20kDa

Immunohistochemistry-Paraffin: Thyroid Peroxidase Antibody [NBP2-27544] - Staining of Human Thyroid Gland. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



Publications

Wiseman SM, Memarnejadian A, Boyce GK et al. Subcutaneous transplantation of human thyroid tissue into a pre-vascularized Cell Pouch(TM) device in a Mus musculus model: Evidence of viability and function for thyroid transplantation PloS one 2022-01-20 [PMID: 35051203] (IF/IHC, Human)

Chen CR, Hamidi S, Braley-Mullen H et al. Antibodies to thyroid peroxidase arise spontaneously with age in NOD.H-2h4 mice and appear after thyroglobulin antibodies. Endocrinology 2010-09-01 [PMID: 20573721]



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA

Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
novus@novusbio.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada

Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: technical@novusbio.com
Orders: orders@novusbio.com
General: novus@novusbio.com

Products Related to NBP2-27544

HAF017	Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
HAF109	Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
NB410-28088-1mg	Goat IgG Isotype Control
H00007173-P01-2ug	Recombinant Human Thyroid Peroxidase GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-27544

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



