Product Datasheet

Human Breast Tissue Lysate (Adult Normal) NBP2-27927

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-27927

Updated 2/12/2018 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-27927



NBP2-27927

Human Breast Tissue Lysate (Adult Normal)

Human breast hissue Lysate (Addit Normal)	
Product Information	
Unit Size	0.1 mg
Concentration	1 mg/ml
Storage	Store at -80C. Avoid freeze-thaw cycles.
Product Description	
Species	Human
Specificity/Sensitivity	Tissue: Breast Diagnosis:Infiltrating ductal carcinoma Stage:IIIB TNM:T4bN1M0 Grade:2 Sex: Female Age :55 Gross findings: Tumor size 10 cm., ill demarcated. Cut section firm and brown/red.
Immunogen	Clinical Tissue Human Breast Normal Tissue Lysate
Preparation Method	Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The pellet was further extracted with a second buffer to obtain the less soluble protein fraction. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature.
Notes	The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides.
Lysate Type	Tissue
Lysate Tissue	Breast
Lysate Tissue Condition	Normal
Lysate Life Stage	Adult
Product Application Details	
Application Notes	These lysates have not been subjected to denaturing or reducing conditions. This allows the tissue or cell lysate to be used in a variety of applications; to study protein-protein interaction, ligand binding, ELISA, immunoprecipitation, 1D and 2D gel electrophoresis, and Western blotting for the detection of specific protein targets. For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel loading buffer with reducing agents may be required. Buffer requirements for performing protein-protein interaction and ligand binding studies can vary significantly from RIPA buffer and may require modifications. In most cases, tissue lysates in RIPA buffer can be used, directly in standard ELISA



This material has tested negative for HbsAg, HIV 1/2, and HCV. Use

treated as a potentially infectious agent and disposed of appropriately.

UNIVERSAL PRECAUTIONS when handling. Human tissue derivatives must be

and immunoprecipitation assays.

Procedures

Product Handling Protocol (NBP2-27927)

Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The pellet was further extracted with a second buffer to obtain the less soluble protein fraction. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature.

Extraction 1: PBS, pH 7.4 1 ug/ml Aprotinin 1 mM NaF

Modified RIPA Buffer: 1 mM EDTA 1 ug/ml Pepstatin-A 0.1% SDS

0.25% Na deoxycholate 1 ug/ml Leupeptin 1 mM PMSF

1 mM Na3VO4

Extraction 2: PBS, pH 7.4, 5.0 M Urea, 2.0 M Thiourea, 50mM DTT, 0.1% SDS





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966 novus@novusbio.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: technical@novusbio.com

Orders: orders@novusbio.com General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-27927

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

