

Product Datasheet

Human Breast Tissue Lysate (Adult Normal) NBP2-27950

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-27950

Updated 2/12/2018 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-27950



NBP2-27950

Human Breast Tissue Lysate (Adult Normal)

Product Information	
Unit Size	0.1 mg
Concentration	1 mg/ml
Storage	Store at -80C. Avoid freeze-thaw cycles.
Product Description	
Species	Human
Specificity/Sensitivity	Tissue: Breast Diagnosis: Infiltrating ductal carcinoma Stage: IV T4N0M0 Grade: 2 Sex: Female Age : 75 Gross findings: Tumor size 8 x 8 cm., ill demarcated. Cut section firm.
Immunogen	Clinical Tissue Human Breast Normal Tissue Lysate
Preparation Method	Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The pellet was further extracted with a second buffer to obtain the less soluble protein fraction. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature.
Notes	The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides.
Lysate Type	Tissue
Lysate Tissue	Breast
Lysate Tissue Condition	Normal
Lysate Life Stage	Adult
Product Application Details	
Application Notes	<p>These lysates have not been subjected to denaturing or reducing conditions. This allows the tissue or cell lysate to be used in a variety of applications; to study protein-protein interaction, ligand binding, ELISA, immunoprecipitation, 1D and 2D gel electrophoresis, and Western blotting for the detection of specific protein targets. For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel loading buffer with reducing agents may be required.</p> <p>Buffer requirements for performing protein-protein interaction and ligand binding studies can vary significantly from RIPA buffer and may require modifications. In most cases, tissue lysates in RIPA buffer can be used, directly in standard ELISA and immunoprecipitation assays.</p> <p>This material has tested negative for HbsAg, HIV 1/2, and HCV. Use UNIVERSAL PRECAUTIONS when handling. Human tissue derivatives must be treated as a potentially infectious agent and disposed of appropriately.</p>

Procedures

Product Handling Protocol (NBP2-27950)

Lysate Preparation

Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The pellet was further extracted with a second buffer to obtain the less soluble protein fraction. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature.

Extraction 1: PBS, pH 7.4 1 ug/ml Aprotinin 1 mM NaF

Modified RIPA Buffer: 1 mM EDTA 1 ug/ml Pepstatin-A 0.1% SDS
0.25% Na deoxycholate 1 ug/ml Leupeptin 1 mM PMSF
1 mM Na₃VO₄

Extraction 2: PBS, pH 7.4, 5.0 M Urea, 2.0 M Thiourea, 50mM DTT, 0.1% SDS





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA

Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
novus@novusbio.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada

Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: technical@novusbio.com
Orders: orders@novusbio.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-27950

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

