Product Datasheet

Human Ovary Tissue Lysate (Adult Tumor) NBP2-28487

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-28487

Updated 2/5/2017 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-28487



NBP2-28487

Human Ovary Tissue Lysate (Adult Tumor)

Product Information		
Concentration 1 mg/ml Storage Store at -80C. Avoid freeze-thaw cycles. Product Description Species Human Specificity/Sensitivity Sex: Female Age :41 Diagnosis: Ovarian papillary adenocarcinoma, moderately differentiated Immunogen Clinical Tissue Human Ovary Tumor Tissue Lysate Preparation Method Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The pellet was further extracted with a second buffer to obtain the less soluble protein fraction. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature. Notes The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. Lysate Type Tissue Lysate Tissue Ovary Lysate Tissue Condition Tumor Lysate Life Stage Adult Product Application Details Application Notes These lysates are proteomic discovery tools.Researchers should validate and optimize for individual use. Potential applications MAY include WB, immunoprecipitation, protein-protein interactions, ligand binding, ELISA. Note: For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel	Product Information	
Storage Store at -80C. Avoid freeze-thaw cycles. Product Description Species Human Sex: Female Age :41 Diagnosis: Ovarian papillary adenocarcinoma, moderately differentiated Immunogen Clinical Tissue Human Ovary Tumor Tissue Lysate Preparation Method Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The pellet was further extracted with a second buffer to obtain the less soluble protein fraction. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature. Notes The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. Lysate Type Tissue Lysate Tissue Ovary Lysate Tissue Condition Tumor Lysate Life Stage Adult Product Application Details Application Notes These lysates are proteomic discovery tools.Researchers should validate and optimize for individual use. Potential applications MAY include WB, immunoprecipitation, protein-protein interactions, ligand binding, ELISA. Note: For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel	Unit Size	0.1 mg
Product Description Species Human Specificity/Sensitivity Sex: Female Age :41 Diagnosis: Ovarian papillary adenocarcinoma, moderately differentiated Immunogen Clinical Tissue Human Ovary Tumor Tissue Lysate Preparation Method Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The pellet was further extracted with a second buffer to obtain the less soluble protein fraction. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature. Notes The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. Lysate Type Tissue Lysate Tissue Ovary Lysate Tissue Condition Lysate Life Stage Adult Product Application Details Application Notes These lysates are proteomic discovery tools. Researchers should validate and optimize for individual use. Potential applications MAY include WB, immunoprecipitation, protein-protein interactions, ligand binding, ELISA. Note: For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel	Concentration	1 mg/ml
Specificity/Sensitivity Sex: Female Age :41 Diagnosis: Ovarian papillary adenocarcinoma, moderately differentiated Immunogen Clinical Tissue Human Ovary Tumor Tissue Lysate Preparation Method Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The pellet was further extracted with a second buffer to obtain the less soluble protein fraction. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature. Notes The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. Lysate Type Tissue Lysate Tissue Ovary Lysate Tissue Condition Lysate Life Stage Adult Product Application Details Application Notes These lysates are proteomic discovery tools. Researchers should validate and optimize for individual use. Potential applications MAY include WB, immunoprecipitation, protein-protein interactions, ligand binding, ELISA. Note: For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel	Storage	Store at -80C. Avoid freeze-thaw cycles.
Specificity/Sensitivity Sex: Female Age :41 Diagnosis: Ovarian papillary adenocarcinoma, moderately differentiated Immunogen Clinical Tissue Human Ovary Tumor Tissue Lysate Preparation Method Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The pellet was further extracted with a second buffer to obtain the less soluble protein fraction. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature. Notes The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. Lysate Type Tissue Lysate Tissue Covary Lysate Tissue Condition Lysate Life Stage Adult Product Application Details Application Notes These lysates are proteomic discovery tools. Researchers should validate and optimize for individual use. Potential applications MAY include WB, immunoprecipitation, protein-protein interactions, ligand binding, ELISA. Note: For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel	Product Description	
Age :41 Diagnosis: Ovarian papillary adenocarcinoma, moderately differentiated Immunogen Clinical Tissue Human Ovary Tumor Tissue Lysate Preparation Method Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The pellet was further extracted with a second buffer to obtain the less soluble protein fraction. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature. Notes The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. Lysate Type Tissue Lysate Tissue Ovary Lysate Tissue Condition Tumor Lysate Life Stage Adult Product Application Details Application Notes These lysates are proteomic discovery tools.Researchers should validate and optimize for individual use. Potential applications MAY include WB, immunoprecipitation, protein-protein interactions, ligand binding, ELISA. Note: For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel	Species	Human
Human Ovary Tumor Tissue Lysate Preparation Method Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The pellet was further extracted with a second buffer to obtain the less soluble protein fraction. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature. Notes The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. Lysate Type Tissue Lysate Tissue Ovary Lysate Tissue Condition Tumor Lysate Life Stage Adult Product Application Details Application Notes These lysates are proteomic discovery tools. Researchers should validate and optimize for individual use. Potential applications MAY include WB, immunoprecipitation, protein-protein interactions, ligand binding, ELISA. Note: For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel	Specificity/Sensitivity	Age :41
proteins, and centrifuged to clarify. The pellet was further extracted with a second buffer to obtain the less soluble protein fraction. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature. Notes The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. Lysate Type Tissue Ovary Lysate Tissue Condition Tumor Lysate Life Stage Adult Product Application Details Application Notes These lysates are proteomic discovery tools. Researchers should validate and optimize for individual use. Potential applications MAY include WB, immunoprecipitation, protein-protein interactions, ligand binding, ELISA. Note: For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel	Immunogen	
Lysate Type Lysate Tissue Ovary Lysate Tissue Condition Lysate Life Stage Adult Product Application Details Application Notes These lysates are proteomic discovery tools.Researchers should validate and optimize for individual use. Potential applications MAY include WB, immunoprecipitation, protein-protein interactions, ligand binding, ELISA. Note: For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel	Preparation Method	proteins, and centrifuged to clarify. The pellet was further extracted with a second buffer to obtain the less soluble protein fraction. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The
Lysate Tissue Condition Lysate Life Stage Adult Product Application Details Application Notes These lysates are proteomic discovery tools. Researchers should validate and optimize for individual use. Potential applications MAY include WB, immunoprecipitation, protein-protein interactions, ligand binding, ELISA. Note: For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel	Notes	
Lysate Tissue Condition Lysate Life Stage Adult Product Application Details Application Notes These lysates are proteomic discovery tools.Researchers should validate and optimize for individual use. Potential applications MAY include WB, immunoprecipitation, protein-protein interactions, ligand binding, ELISA. Note: For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel	Lysate Type	Tissue
Lysate Life Stage Adult Product Application Details Application Notes These lysates are proteomic discovery tools.Researchers should validate and optimize for individual use. Potential applications MAY include WB, immunoprecipitation, protein-protein interactions, ligand binding, ELISA. Note: For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel	Lysate Tissue	Ovary
Product Application Details Application Notes These lysates are proteomic discovery tools.Researchers should validate and optimize for individual use. Potential applications MAY include WB, immunoprecipitation, protein-protein interactions, ligand binding, ELISA. Note: For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel	Lysate Tissue Condition	Tumor
Application Notes These lysates are proteomic discovery tools.Researchers should validate and optimize for individual use. Potential applications MAY include WB, immunoprecipitation, protein-protein interactions, ligand binding, ELISA. Note: For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel	Lysate Life Stage	Adult
optimize for individual use. Potential applications MAY include WB, immunoprecipitation, protein-protein interactions, ligand binding, ELISA. Note: For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel	Product Application Details	
	Application Notes	optimize for individual use. Potential applications MAY include WB, immunoprecipitation, protein-protein interactions, ligand binding, ELISA. Note: For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel



Procedures

Product Handling Protocol (NBP2-28487)

Lysate Preparation:

Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The pellet was further extracted with a second buffer to obtain the less soluble protein fraction. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature.

Extraction 1: PBS, pH 7.4, 1 ug/ml Aprotinin, 1 mM NaF

Modified RIPA Buffer: 1 mM EDTA, 1 ug/ml Pepstatin-A, 0.1% SDS, 0.25% Na deoxycholate, 1 ug/ml Leupeptin, 1 mM PMSF, 1 mM Na3VO4

Extraction 2: PBS, pH 7.4, 5.0 M Urea, 2.0 M Thiourea, 50mM DTT, 0.1% SDS





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966 novus@novusbio.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: technical@novusbio.com

Orders: orders@novusbio.com General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-28487

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

