

Product Datasheet

Human Ovary Tissue Lysate (Adult Tumor) NBP2-28495

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-28495

Updated 2/5/2017 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-28495



NBP2-28495**Human Ovary Tissue Lysate (Adult Tumor)**

| Product Information | |
|------------------------------------|--|
| Unit Size | 0.1 mg |
| Concentration | 1 mg/ml |
| Storage | Store at -80C. Avoid freeze-thaw cycles. |
| Product Description | |
| Species | Human |
| Specificity/Sensitivity | Sex: Female Age :50 Diagnosis: Ovarian papillary cystadenocarcinoma, moderately differentiated |
| Immunogen | Clinical Tissue Human Ovary Tumor Tissue Lysate |
| Preparation Method | Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The pellet was further extracted with a second buffer to obtain the less soluble protein fraction. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature. |
| Notes | The vial is provided with a 10% overfill. Maximum recovery can be obtained by centrifuging the vial briefly to collect any solution on the cap and tube sides. |
| Lysate Type | Tissue |
| Lysate Tissue | Ovary |
| Lysate Tissue Condition | Tumor |
| Lysate Life Stage | Adult |
| Product Application Details | |
| Application Notes | These lysates are proteomic discovery tools. Researchers should validate and optimize for individual use. Potential applications MAY include WB, immunoprecipitation, protein-protein interactions, ligand binding, ELISA. Note: For use in 1D and 2D gel electrophoresis, the addition of a denaturing gel loading buffer with reducing agents may be required. |

Procedures

Product Handling Protocol (NBP2-28495)

Lysate Preparation:

Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The pellet was further extracted with a second buffer to obtain the less soluble protein fraction. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature.

Extraction 1: PBS, pH 7.4, 1 ug/ml Aprotinin, 1 mM NaF

Modified RIPA Buffer: 1 mM EDTA, 1 ug/ml Pepstatin-A, 0.1% SDS, 0.25% Na deoxycholate, 1 ug/ml Leupeptin, 1 mM PMSF, 1 mM Na₃VO₄

Extraction 2: PBS, pH 7.4, 5.0 M Urea, 2.0 M Thiourea, 50mM DTT, 0.1% SDS





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA

Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
novus@novusbio.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada

Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: technical@novusbio.com
Orders: orders@novusbio.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-28495

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

