Product Datasheet

BST2 Antibody (2E2) - BSA Free NBP2-29621

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-29621

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.


Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-29621




| BST2 Antibody (2E2) - BSA Free | |
|--------------------------------|---|
| Product Information | |
| Unit Size | 0.1 mg |
| Concentration | 1.0 mg/ml |
| Storage | Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles. |
| Clonality | Monoclonal |
| Clone | 2E2 |
| Preservative | 0.05% Sodium Azide |
| Isotype | IgG1 Kappa |
| Purity | Protein A or G purified |
| Buffer | PBS |
| Product Description | |
| Host | Mouse |
| Gene ID | 684 |
| Gene Symbol | BST2 |
| Species | Human |
| Immunogen | Full length recombinant human BST2 protein [NCBI NP_004326] |
| Product Application Details | |
| Applications | Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Paraffin, CyTOF-ready |
| Recommended Dilutions | Western Blot 1 ug/ml, Flow Cytometry 0.5 ug/ml, Immunohistochemistry 5 ug/ml, Immunohistochemistry-Paraffin 5 ug/ml, CyTOF-ready |
| Application Notes | The unprocessed form of BST2 protein is 180 amino acids long (161 amino acids in its cleaved BST) and its cleaved form undergoes further modifications namely disulfide bond formation, GPI-anchor/GPI-like-anchor binding, glycosylation, isopeptide bond formation, lipidation and UbI conjugation. The predicted molecular weight of unmodified BST2 protein is ~17-18kDa, whereas the post-translationally modified protein has been documented to run at ~25-36kDa (glycosylated), 50-55 kDa (ubiquitinated) and 65-70 kDa (dimers) in WB assay. BST2 protein localize mainly to the cell membranes from where it shuttles to the trans-Golgi network, late endosome and the cytoplasm. In immuno-staining assays, BST2 may develop a membrane-cytoplasmic staining pattern. This antibody is CyTOF ready. |

Images


Western Blot: BST2 Antibody (2E2) [NBP2-29621] - Analysis of BST2 protein in (A) human heart lysate, (B) human ovary lysate and (C) partial recombinant BST2 protein with monoclonal BST2 antibody at a concentration of 1 ug/ml. In the tested human lysates, two distinct bands were observed which represents the glycosylated (lower bands) and the ubiquitinated (higher bands) forms of BST2 protein.




Immunohistochemistry-Paraffin: BST2 Antibody (2E2) [NBP2-29621] - Analysis of BST2 protein in a section of human endometrial carcinoma using 5 ug/ml concentration of BST2 antibody (clone 2E2). The carcinoma cells depicted distinct membrane-cytoplasmic BST2 positivity with more intense staining in the cellular membranes, whereas, weak staining was observed in tumor stroma.



Immunohistochemistry-Paraffin: BST2 Antibody (2E2) [NBP2-29621] - Analysis of BST2 protein in a section of human renal cell carcinoma (clear cell type) using 5 ug/ml concentration of BST2 antibody (clone 2E2). The carcinoma cells depicted distinct membrane-cytoplasmic BST2 positivity with more intense staining in cellular membranes.



Immunohistochemistry-Paraffin: BST2 Antibody (2E2) [NBP2-29621] - Analysis of BST2 protein in a section of human lung cancer (squamous cell carcinoma) using 5 ug/ml concentration of BST2 antibody (clone 2E2). The carcinoma cells depicted distinct membrane-cytoplasmic BST2 positivity with more intense staining in the cellular membranes, whereas, mild to moderate staining was observed in tumor stroma also.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-29621

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-43319-0.5mg Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)

NBP2-29621AF405 BST2 Antibody (2E2) [Alexa Fluor® 405]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-29621

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

