Product Datasheet

INSTA-Blot Breast Tissue OncoPair NBP2-30126

Unit Size: 1 Membrane

Store at room temperature.

www.novusbio.com



technical@novusbio.com

Publications: 5

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-30126

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-30126



NBP2-30126

Product Information

INSTA-Blot Breast Tissue OncoPair

Product Information	
Unit Size	1 Membrane
Concentration	Concentration is not relevant for this product. Please see the protocols for proper use of this product.
Storage	Store at room temperature.
Product Description	
Description	Please see online datasheet for lane details: www.novusbio.com/NBP2-30126
Preparation Method	Tissue specimens were homogenized in modified RIPA buffer to obtain the soluble proteins and centrifuged to clarify. The concentration of protein in each lysate was determined by a standard protein assay and standardized to 1 mg/ml. Sample buffer was added to the soluble fraction and approximately 14 ug/lane of protein was run and then transferred to PVDF membraneSoluble fraction extraction buffer (extract 1): PBS at pH 7.4, 1 ug/ml Aprotinin, 1 mM NaF - Modified RIPA buffer: 1 mM EDTA, 1 ug/ml Pepstatin-A, 0.1% SDS , 0.25% Na deoxycholate, 1 ug/ml Leupeptin, 1 mM PMSF, 1 mM Na3VO4 -5X Sample buffer: 50 ml glycerol, 15 g SDS, 3.819 g Tris, pH 6.8, 25 ml 2-ME, 100 mg bromophenol blue, final volume of 100 ml with DI H2O.

Publications

Santagata S, Hu R, Lin NU et al. High levels of nuclear heat-shock factor 1 (HSF1) are associated with poor prognosis in breast cancer. Proc Natl Acad Sci U S A. 2011-11-08 [PMID: 22042860]

Details:

INSTA-BLOT(TM) Breast Tissue OncoPair (IMB-130a). Membrane was probed with HSF1 antibody, Fig 1C.

Tuszynski GP, Rothman VL, Zheng X et al. G-protein coupled receptor-associated sorting protein 1 (GASP-1), a potential biomarker in breast cancer. Exp Mol Pathol. 2011-10-01 [PMID: 21791203]

Details:

INSTA-BLOT(TM) Breast Tissue OncoPair (IMB-130e). Blot was probed with a GASP-1 antibody, Fig 2.2. Breast cancer Tissue Microarray (IMH-364, aka CBA4). Tissue microarray was probed with a GASP-1 antibody, Figs 3, 4.

Qin Y, Xu J, Aysola K et al. Ubc9 mediates nuclear localization and growth suppression of BRCA1 and BRCA1a proteins. J Cell Physiol. 2011-12-01 [PMID: 21344391]

Details:

INSTA-BLOT(TM) Breast Tissue OncoPair (IMB-130e). Membrane was probed with Ubc9 antibody, Fig 7A.

Michel V, Yuan Z, Ramsubir S et al. Choline transport for phospholipid synthesis. Exp Biol Med (Maywood). 2006-05-01 [PMID: 16636297] (WB, Human)

Details:

Product cited: INSTA-Blot(TM) Human Tissue: ready-to-use WB (IMB-103).

Kinouchi R, Kinouchi T, Hamamoto T et al. Distribution of CESP-1 protein in the corneal endothelium and other tissues. Invest Ophthalmol Vis Sci. 2006-04-01 [PMID: 16565373] (WB, Human)

Details:

INSTA-BLOT(TM) Human Tissue: ready-to-use WB (IMB-103).



Procedures

Western Blot Protocol for INSTA-Blot Breast Tissue OncoPair (NBP2-30126)

Note: The INSTA-Blot(TM) PVDF membrane has been dried and must be rehydrated (Step one) prior to use.

Before wetting, it is suggested that the top of the lanes be marked with an ink pen that will not wash off in methanol.

- 1. Wet the blots with 100% methanol then thoroughly wash with TBST (25 mM Tris-Cl, pH 8.0; 125 mM NaCl; 0.1% Tween 20) twice to remove residual methanol.
- 2. Incubate the blot for 1 hr with 5% Carnation nonfat dry milk in TBST to block non-specific antibody binding.
- 3. Incubate the blots with primary antibody in 1% milk/TBST for 1-2 h at RT or overnight at 4 degrees C.
- 4. After incubation with the primary antibody, wash the blots five times in TBST then incubate with a secondary antibody conjugated to horseradish peroxidase (HRP-conjugated secondary antibodies) for 1-2 h at RT.
- 5. After five washes with TBST, develop the blots for 5 min using the PicoTect(TM) Western Blot Chemiluminescent Substrate.
- 6. Expose the blots to photographic film for an appropriate time period. We normally use Hyper-film(TM)-ECL films (Amersham Life Science Inc.) and expose for various periods ranging from 10 s to 20 min to visualize the chemiluminescence signal corresponding to the specific antibody-antigen reaction.



MSDS for Non Hazardous Product (NBP2-30126)

Material Safety Data Sheet for Non Hazardous Products

Hazard Information

Chemical Name: Non hazardous products.

Chemical Formula: N/A CAS Number: N/A EEC-No: N/A

Hazard Identification

None

First Aid Measures Eye Contact: None Skin Contact: None Inhalation: None Ingestion: None

Accidental Release Measures

This product either does not contain hazardous constituents or the concentration of all chemical constituents are below the regulatory threshold limits described by Occupational Safety Health Administration Hazard Communication Standard 29 CFR 1910.1200 and the European Directive 91/155/EEC. 88/379/EEC, and 67/546/EEC.

Handling and Storage

Exposure Controls / Personal Protection

Other Precautions: None

Physical and Chemical Properties

Form: N/A Color: N/A Odor: N/A

Melting Point: N/A

Boiling Temperature: N/A

Density: N/A

Vapor Pressure: N/A Solubility in Water: N/A Flash Point: N/A Explosion limits: N/A Ignition Temperature: N/A





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@bio-techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Western Blot Membranes are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-30126

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

