Product Datasheet

TIRAP (TLR2 and TLR4) Inhibitor Peptide Set NBP2-31274-1mg

Unit Size: 1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-31274

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-31274



NBP2-31274-1mg

TIRAP (TLR2 and TLR4) Inhibitor Peptide Set

Product Information	
Unit Size	1 mg
Concentration	Lyoph
Storage	Store at -20C. Avoid freeze-thaw cycles.
Reconstitution Instructions	Please contact technical support for detailed reconstitution instructions.
Buffer	Lyophilized white powder
Product Description	
Gene ID	114609
Gene Symbol	TIRAP
Species	Human, Mouse
Specificity/Sensitivity	TLR2 and TLR4 Inhibitor Peptide: COBRA TLR2 and TLR4 Inhibitor interferes with interaction between TIRAP/Mal and TIR domain of TLR2 or TLR4.
Immunogen	TLR2 / TLR4 Inhibitor Peptide: 1 mg (lyophilized); sequence: DRQIKIWFQNRRMKWKKPGFLRDPWCKYQML (Inhibitor sequence: PGFLRDPWCKYQML);
Preparation Method	 Preparation of 2.5 mM peptide stock solutions TLR2 / TLR4 Inhibitor Peptide: A final volume of 100 ul will make a 2.5 mM stock solution. Add 100 ul sterile water to the tube of peptide. Carefully pipet to ensure all of the peptide is dissolved and briefly spin the tube before opening. The stock solutions may be diluted further to make working solutions. Dilute according to the needs for your assay. For example, dilute 2.5 mM stock solutions 1:10 in sterile 1X PBS or cell culture media to make 250 uM working solutions. Working solution should be made fresh daily and not be stored.
Inhibitor Family	TLR
Inhibitor Target	TLR2 and TLR4
Inhibitor Content	Antennapedia Control Peptide: 1 mg (lyophilized); sequence: DRQIKIWFQNRRMKWKK; Molecular weight: 2361 Da
Product Application Details	
Applications	Flow Cytometry, Functional
Recommended Dilutions	Flow Cytometry, Functional
Application Notes	The inhibitor is used in assays to inhibit TLR2 and TLR4 signaling. We recommend an initial titration of the inhibitor from 0-50 uM for in vitro assays along with control of which concentrations should be mirror inhibitor concentrations. Inhibitor and control should be preincubated with cells prior to ligand activation to allow sufficient time for the peptides to enter from the media into the cell. We typically preincubate with inhibitor and control for 1 h prior to TLR2 or TLR4 activation (Figures 1 and 2); however, optimal preincubation times may vary between model systems. ELOW and Eunctional application reported by

may vary between model systems. FLOW and Functional application reported by Alvarez et al - Conference on Retroviruses and Opportunistic Infections - CROI 2014 (Abstract: The NCOR2-Nurr1-CoREST Transrepression Axis Impairs HIV Reactivation in Latently Infected Microglial Cells)



Publications

Alvarez D, Das B, Garcia-Mesa Y et al. The NCOR2-Nurr1-CoREST Transrepression Axis Impairs HIV Reactivation in Latently Infected Microglial Cells Conference on Retroviruses and Opportunistic Infections (CROI) 2014. (Func, FLOW, Human)

Details:

TIRAP/COBRA (TLR2 or TLR4) inhibitor used at 20uM concentration on Microglial/HIV Cells - CHME-5/HIV and hT-HMucroglia/HIV cells were untreated or treated with IL-1 beta (100 pg/mL), Pam3CSK4 (0.1 ug/mL), or a combination of IL-1 beta and Pam3CSK4 for 16 h prior to measuring GFP-expressing cells by FLOW (Fig 4B). COBRA, but not of VIPER, abolished the effect of Pam3CSK4, but not of IL-1 beta indicating that Pam3CSK4 is indeed responsible for the potentiation of HIV reactivation





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP2-31274-1mg

M6000B-1	IL-6 [HRP]
NBP2-08583	TIRAP (TLR2 and TLR4) Overexpression Lysate
210-TA-005	TNF-alpha [Unconjugated]
NBP2-22871	Recombinant Human TIRAP (TLR2 and TLR4) His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Inhibitors are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-31274

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

