

Product Datasheet

NALP4 Antibody - BSA Free

NBP2-31375

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-31375

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-31375



NBP2-31375

NALP4 Antibody - BSA Free

Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	PBS


Product Description	
Description	Novus Biologicals Rabbit NALP4 Antibody - BSA Free (NBP2-31375) is a polyclonal antibody validated for use in WB and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	147945
Gene Symbol	NLRP4
Species	Human
Reactivity Notes	Immunogen shows 51% similarity with Mouse's NALP4.
Specificity/Sensitivity	Besides detecting canonical isoform (~113.4 kDa), this antibody may detect isoform 3 (104 kDa) also.
Immunogen	Partial recombinant human NALP4 protein (between residues 1-300) [UniProt Q96MN2]

Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence
Recommended Dilutions	Western Blot 1-2 ug/ml, Immunocytochemistry/ Immunofluorescence 0.1 ug/ml




Images

Western Blot: NALP4 Antibody [NBP2-31375] - WB detection of NALP4/ NLRP4 protein in (A) K562 cell lysate and (B) partial recombinant protein by using NALP4 antibody at a concentration of: 1 ug/ml for lysate and 0.05 ug/ml for recombinant protein. In K562 cells lysate, this antibody detected a single specific band at ~116 kDa position (predicted size of NALP4 is 113.4 kDa).



Immunocytochemistry/Immunofluorescence: NALP4 Antibody [NBP2-31375] - NALP4 antibody was tested in MCF7 cells with Dylight 488 (green). Nuclei and alpha-tubulin were counterstained with DAPI (blue) and Dylight 550 (red). An antibody concentration of 0.1 ug/ml was used. Image objective 40x.



Procedures

Western Blot protocol for NALP4 Antibody (NBP2-31375)

NALP4 Antibody:

Reagents needed:

- a. Washing Buffer: Tris Buffer Saline with 0.01% of tween 20).
- b. Blocking Buffer: 5% skimmed milk powder in washing buffer).
- c. Secondary antibody, Horseradish peroxidase conjugated.
- d. Chemiluminescent solution (SuperSignal WestPico™, Pierce).

Western blot Method:

1. Perform SDS-PAGE using PVDF membrane. Cut into strips.
2. Activate strips with methanol by dipping them into methanol for 5 min.
3. Discard the methanol and take fresh methanol to repeat step b.
4. Let the strips dry, and then add blocking solution and incubate at RT in a shaker for 30-45 minutes.
5. Dilute primary antibody in blocking buffer. Incubate the number of strips required with the diluted primary antibody at room temperature for 2 hours in a shaker.
6. Wash strips two times with washing buffer at 30 minutes intervals.
7. Dilute HRP conjugated secondary antibody in blocking buffer. Add diluted secondary antibody to the membrane strips and incubate for exactly 1 hour while shaking at RT.
8. Wash the strips with washing buffer for 2-3 hours with 3 to 4 changes on a shaker. This helps in reducing the background staining.
9. Prepare the chemiluminescent solution (SuperSignal WestPico™) by mixing solution A and Solution B at 1:1. Mix well. Soak the strip in the chemiluminescent solution; keep for 3-5 minutes under constant shaking.
10. Expose the membrane to a sheet of film and develop.

Immunocytochemistry/Immunofluorescence protocol for NALP4 Antibody (NBP2-31375)

NALP4 Antibody:

Immunocytochemistry Protocol

Culture cells to appropriate density on suitable glass coverslips in 35 mm culture dishes or 6-well plates.

1. Remove culture medium and add 10% formalin to the dish. Fix at room temperature for 5-10 minutes.
2. Remove the formalin and add 0.5% Triton-X 100 in TBS to permeabilize the cells. Incubate for 5-10 minutes.
3. Remove the permeabilization buffer and add wash buffer (i.e. PBS or PBS with 0.1% Tween-20). Be sure to not let the specimen dry out. Gently wash three times for 10 minutes.
4. Alternatively, cells can be fixed with -20C methanol for 10 min at room temperature. Remove the methanol and rehydrate in PBS for 10 min before proceeding.
5. To block nonspecific antibody binding incubate in 10% normal goat serum for 1 hour at room temperature.
6. Add primary antibody at appropriate dilution and incubate at room temperature for 1 hour or at 4 degrees C overnight.
7. Remove primary antibody and replace with wash buffer. Gently wash three times for 10 minutes.
8. Add secondary antibody at the appropriate dilution. Incubate for 1 hour at room temperature.
9. Remove antibody and replace with wash buffer. Gently wash three times for 10 minutes.
10. Nuclei can be staining with 4',6' diamino phenylindole (DAPI) at 0.1 ug/ml, or coverslips can be directly mounted in media containing DAPI.
11. Cells can now be viewed with a fluorescence microscope.

*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow proper laboratory procedures for the disposal of formalin.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-31375

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NB100-56156PEP	NALP4 Antibody Blocking Peptide

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-31375

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

