# **Product Datasheet**

# FoxO1/FKHR Antibody (83N7F8) - BSA Free NBP2-31376SS

Unit Size: 0.025 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 1 Publications: 7

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-31376

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-31376



# NBP2-31376SS


FoxO1/FKHR Antibody (83N7F8) - BSA Free	
Product Information	
Unit Size	0.025 mg
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	83N7F8
Preservative	0.05% Sodium Azide
Isotype	IgG2b Kappa
Purity	Protein G purified
Buffer	PBS
Product Description	
Description	Novus Biologicals Mouse FoxO1/FKHR Antibody (83N7F8) - BSA Free (NBP2-31376) is a monoclonal antibody validated for use in IHC, WB, Flow, ICC/IF and Simple Western. Anti-FoxO1/FKHR Antibody: Cited in 7 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	2308
Gene Symbol	FOXO1
Species	Human, Mouse, Rat
Reactivity Notes	Use in Mouse reported in scientific literature (PMID:33662610). Rat reactivity reported in scientific literature (PMID: 31553623).
Immunogen	FOXO1A
Product Application Details	
Applications	Western Blot, Simple Western, Immunohistochemistry-Paraffin, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 0.5-2 ug/ml, Simple Western 1:50, Flow Cytometry, Immunohistochemistry 5 ug/ml, Immunocytochemistry/ Immunofluorescence 0.1-2 ug/ml, Immunohistochemistry-Paraffin 5 ug/ml
Application Notes	In Simple Western only 10 - 15 uL of the recommended dilution is used per data point.  See <u>Simple Western Antibody Database</u> for Simple Western validation: Tested in Hek293 lysate 0.5 mg/mL, separated by Size, antibody dilution of 1:50, apparent MW was 107 kDa




MW was 107 kDa.

## **Images**


Western Blot: FoxO1/FKHR Antibody (83N7F8) [NBP2-31376] - Analysis of (A) Ntera2 cell lysate, (B) HEK293 cell lysate, and (C) human placenta tissue using FOXO1A antibody at a concentration of 2 ug/ml.




Immunocytochemistry/Immunofluorescence: FoxO1/FKHR Antibody (83N7F8) [NBP2-31376] - FOXO1A antibody was tested in Ntera2 cells with DyLight 488 (green). Nuclei and alpha-tubulin were counterstained with DAPI (blue) and DyLight 550 (red). An antibody concentration of 0.1 ug/ml was used. Image objective 40x.




Immunohistochemistry-Paraffin: FoxO1/FKHR Antibody (83N7F8) [NBP2 -31376] - Analysis of FOXO1A protein in a section of normal human uterus by using FOXO1A antibody at a concentration of 5 ug/ml. The columnar cells of straight non-convoluted glands in uterous showed specific staining for FOXO1A in the cellular cytoplasm as well as nuclei, whereas the stromal cells showed mild cytoplasmic staining only.



Flow Cytometry: FoxO1/FKHR Antibody (83N7F8) [NBP2-31376] - An intracellular stain was performed on RH-30 cells with FoxO1/FKHR Antibody [83N7F8] NBP2-31376AF488 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 488.




Western Blot: FoxO1/FKHR Antibody (83N7F8) [NBP2-31376] Detection of FOXO1A / FOXO1 protein in (A) Hela cells lysate, (B) Jurkat
cell lysate, and (C) partial recombinant protein by using FOXO1A
antibody at a concentration of 0.5 ug/ml for lysate and 0.05 ug/ml for the
recombinant protein.



Immunohistochemistry-Paraffin: FoxO1/FKHR Antibody (83N7F8) [NBP2 -31376] - Analysis of FOXO1A protein in a section of normal human kidney by using FOXO1A antibody at a concentration of 5 ug/ml. Cytoplasmic and nuclear positivity was observed in the cells of various tubules/ducts of renal medulla.




Flow Cytometry: FoxO1/FKHR Antibody (83N7F8) [NBP2-31376] - Analysis using the Alexa Fluor 488 conjugated FoxO1 antibody, NBP2-31376AF488 (blue) and fluorescence minus one control (red). Staining of FoxO1 in primary CD4+ T-cells isolated from human peripheral blood using 1uL of antibody per 1X10e5 cells. Image from verified customer review.




CD4+ T-cells derived from human peripheral blood were stained with Alexa Fluor 488 conjugated anti-FOXO1 antibody (blue) from Novus Biologicals (Littleton, CO; Catalog # NBP2-31376AF488). Fluorescence minus one control is shown in red.

Flow Cytometry: FoxO1/FKHR Antibody (83N7F8) [NBP2-31376] - Using the Alexa Fluor 488 direct conjugate An intracellular stain was performed on Daudi cells with FoxO1/FKHR (83N7F8) antibody NBP2-31376AF488 (blue) and a matched isotype control NBP2-27231AF488 (orange). Cells were fixed with 4% PFA and then permeablized with 0.1% saponin. Cells were incubated in an antibody dilution of 5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 488.



Flow Cytometry: FoxO1/FKHR Antibody (83N7F8) [NBP2-31376] - An intracellular stain was performed on U2-OS cells with FoxO1/FKHR [83N7F8] Antibody NBP2-31376AF488 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 10 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 488.



Simple Western: FoxO1/FKHR Antibody (83N7F8) [NBP2-31376] - Lane view shows a specific band for FoxO1/FKHR in 0.5 mg/ml of Hek293 lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.



#### **Publications**

Tan L, Shi G, Zhao J et al. MDSCs participate in the pathogenesis of diffuse pulmonary hemorrhage in murine lupus through mTOR-FoxO1 signaling Biochemistry and Biophysics Reports 2022-12-01 [PMID: 36164563]

Zhang W, Cao X, Zhong X et al. Steroid nuclear receptor coactivator 2 controls immune tolerance by promoting induced Treg differentiation via up-regulating Nr4a2 Science advances 2022-06-17 [PMID: 35704583] (FLOW, Mouse)

Li D, Lu L, Kong W et al. C-type lectin receptor Dectin3 deficiency balances the accumulation and function of FoxO1-mediated LOX-1(+) M-MDSCs in relieving lupus-like symptoms Cell Death Dis 2021-09-03 [PMID: 34480018]

#### Details:

Citation using the Alexa Fluor 750 format of this antibody.

Zhang X, Liu X, Wan F et al. Protective effect of resveratrol against hydrogen peroxide-induced oxidative stress in bovine skeletal muscle cells Meat science 2021-12-16 [PMID: 34952489] (WB)

Ding Z, Qiu M, Alharbi M et al FOXO1 expression in chondrocytes modulates cartilage production and removal in fracture healing Bone 2021-03-04 [PMID: 33662610] (FLOW, Mouse)

#### Details:

Citation using the Alexa Fluor 488 version of this antibody.

Presby DM, Checkley LA, Jackman MR et al. Regular exercise potentiates energetically expensive hepatic de novo lipogenesis during early weight regain Am. J. Physiol. Regul. Integr. Comp. Physiol. 2019-09-25 [PMID: 31553623] (WB, Rat)

Hepp P, Hutter S, Knabl J et al. Histone H3 lysine 9 acetylation is downregulated in GDM Placentas and Calcitriol supplementation enhanced this effect Int J Mol Sci 2018-12-14 [PMID: 30558244] (WB, Human)





# Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

# **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

# **General Contact Information**

www.novusbio.com Technical Support: nb-technical@bio-

Orders: nb-customerservice@bio-techne.com

techne.com

General: novus@novusbio.com

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-31376

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

