

Product Datasheet

Delta Np73 Antibody (38C674.2) - Azide and BSA Free NBP2-33262

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP2-33262

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP2-33262



NBP2-33262

Delta Np73 Antibody (38C674.2) - Azide and BSA Free


Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	38C674.2
Preservative	No Preservative
Isotype	IgG1 Kappa
Purity	Protein G purified
Buffer	PBS

Product Description	
Description	Novus Biologicals Mouse Delta Np73 Antibody (38C674.2) - Azide and BSA Free (NBP2-24873) is a monoclonal antibody validated for use in IHC, WB and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	7161
Gene Symbol	TP73
Species	Human, Mouse, Rat
Immunogen	This antibody was developed against a peptide corresponding to a portion of amino acid residues 1-50 of human dNp73. Mouse and human sequences are more than 95 % identical at these amino acid residues. In mouse, alanine at position 8 is replaced by methionine.


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen
Recommended Dilutions	Western Blot 1-3 ug/ml, Immunohistochemistry, Immunocytochemistry/Immunofluorescence 1:20-1:1000, Immunohistochemistry-Paraffin 5 ug/ml, Immunohistochemistry-Frozen 1:10-1:500
Application Notes	Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM sodium citrate buffer, pH 6.0 for 10-20 min followed by cooling at RT for 20 min.

Images


Western Blot: Delta Np73 Antibody (38C674.2) - Azide Free [NBP2-33262] - Delta Np73 Antibody (38C674.2) analysis for Np73 using NBP2-24873 at 1 ug/ml in A) a cell line transfected with Np73 cDNA, B) HeLa, and C) NIH 3T3 cell lysate. Image from the standard format of this antibody.




Immunocytochemistry/Immunofluorescence: Delta Np73 Antibody (38C674.2) - Azide Free [NBP2-33262] - HeLa cells were fixed for 10 minutes using 10% formalin and then permeabilized for 5 minutes using 1X TBS + 0.5% Triton-X100. The cells were incubated with anti-Delta Np73 (38C674.2) NBP2-24873 at a 1:50 dilution overnight at 4C and detected with an anti-mouse Dylight 488 (Green) at a 1:500 dilution. Actin was counterstained with Phalloidin 568 (Red) at a 1:200 dilution. Nuclei were counterstained with DAPI (Blue). Cells were imaged using a 40X objective. Image from the standard format of this antibody.




Immunohistochemistry-Paraffin: Delta Np73 Antibody (38C674.2) - Azide Free [NBP2-33262] - Delta Np73 Antibody (38C674.2) Formalin-fixed, paraffin-embedded lung squamous cell carcinoma probed with p73 antibody at 5 ug/ml. Image from the standard format of this antibody.



Immunohistochemistry: Delta Np73 Antibody (38C674.2) - Azide Free [NBP2-33262] - Delta Np73 Antibody (38C674.2) analysis of Delta Np73 in embryonic day 15 (A) and day 17 (B) E15 rat embryonic kidney at 1:2000. Delta Np73 is expressed in ureteric bud (UB) branches, and metanephrogenic mesenchyme (MM), as well as in cells invading the g



Immunohistochemistry-Paraffin: Delta Np73 Antibody (38C674.2) - Azide Free [NBP2-33262] - Delta Np73 Antibody (38C674.2) Staining of human placenta. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 5 ug/ml. Image from the standard format of this antibody.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP2-33262

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
NB100-355	RPE65 Antibody (401.8B11.3D9) - BSA Free

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-33262

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

