Product Datasheet

SMC3 [p Ser103] Antibody - BSA Free NBP2-37694

Unit Size: 100 ul

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP2-37694

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP2-37694



NBP2-37694

SMC3 [p Ser103] Antibody - BSA Free


SMC3 [p Ser103] Antibody - BSA Free	
Product Information	
Unit Size	100 ul
Concentration	1.0 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris-Citrate/Phosphate (pH 7.0 - 8.0)
Product Description	
Description	Novus Biologicals Rabbit SMC3 [p Ser103] Antibody - BSA Free (NBP2-37694) is a polyclonal antibody validated for use in WB and IP. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	9126
Gene Symbol	SMC3
Species	Human
Immunogen	This antibody maps to a region between residue This antibody maps to a region surrounding residue 1083 when Serine 1083 is phosphorylated of human Structural Maintenance of Chromosomes 3 using the numbering given in SwissProt entry Q9UQE7 (GeneID 9126). of human Structural Maintenance of Chromosome 3 using the numbering given in entry NP_005436.1 (GeneID 9126).
Product Application Details	
Applications	Western Blot, Immunoprecipitation
Recommended Dilutions	Western Blot 1:500 - 1:2500, Immunoprecipitation 2 - 10 ug/mg lysate
Application Notes	Antibody concentration was determined by extinction coefficient: absorbance at

Product Application Details	
Applications	Western Blot, Immunoprecipitation
Recommended Dilutions	Western Blot 1:500 - 1:2500, Immunoprecipitation 2 - 10 ug/mg lysate
Application Notes	Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.




Images


Western Blot: SMC3 [p Ser103] Antibody [NBP2-37694] - Detection of human Phospho SMC3 (S1083) by western blot (WB). Samples: Whole cell lysate (50 ug) from HeLa cells treated with 100 uM EPE and/or 100 uM phosphatase for 4 hours (+) or mock treated (-) cells. Antibodies: Affinity purified rabbit anti-Phospho SMC3 (S1083) antibody NBP2-37694 used for WB at 1 ug/ml (upper panel); to detect total SMC3, another rabbit anti-SMC3 was used for WB at 0.1 ug/ml (lower panel). Detection: Chemiluminescence with an exposure time of 3 minutes (upper panel) or 10 seconds (lower panel).




Western Blot: SMC3 [p Ser103] Antibody [NBP2-37694] - Whole cell lysate (50 ug) from HeLa cells treated with 500 uM EPE for 4 hours (+) or mock treated (-) cells Affinity purified rabbit anti-Phospho SMC3 (S1083) antibody used for WB at 1 ug/ml (upper panel); to detect total SMC3, rabbit anti-SMC3 was used for WB at 0.1 ug/ml (lower panel). Detection: Chemiluminescence with an exposure time of 3 minutes (upper panel) or 30 seconds (lower panel).



Immunoprecipitation: SMC3 [p Ser103] Antibody [NBP2-37694] - Whole cell lysate (1mg for IP; 20% of IP loaded) from HeLa cells treated with 500 uM EPE for 4 hours (+). Antibodies: Phospho SMC3 was immunoprecipitated (IP) using rabbit anti-Phospho SMC3 (S1083) antibody. The IP reactions were either mock treated (-) or treated with phosphatases (+). Affinity purified rabbit anti-Phospho SMC3 (S1083) antibody was used for western blot (WB) of IPs at 1 ug/ml (upper panel); to detect total SMC3 affinity purified rabbit anti-SMC3 was used for WB of IPs at 0.1 ug/ml (lower panel). Detection: Chemiluminescence with an exposure time of 3 minutes (upper panel) and 30 seconds (lower panel).



SMC3 [p Ser103] Antibody [NBP2-37694] - Detection of human Phospho SMC3 (S1083) by western blot of immunoprecipitates. Samples: Whole cell lysate (1mg for IP; 20% of IP loaded) from HeLa cells treated with 100 uM EPE for 4 hours (+) or mock treated (-). Antibodies: Phospho SMC3 (S1083) was immunoprecipitated using rabbit anti-Phospho SMC3 (S1083) antibody NBP2-37694 at 6 ug per reaction. Total SMC3 was immunoprecipitated with another rabbit anti-SMC3 antibody. Detection: Chemiluminescence with an exposure time of 10 seconds.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP2-37694

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

H00009126-Q01-10ug Recombinant Human SMC3 GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP2-37694

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

